Issue number: BT-NG-020627-560-0032

Proposed Grid Supply Point Substation Off the

Environmental Appraisal Appendix 3: Biodiversity Baseline April 2022

nationalgrid

Page left intentionally blank

Contents

Appe	endix 3: Biodiversity Baseline	1
1.	Introduction	1
1.1	Purpose of this Document	1
1.2	Legislation	1
2.	Methodology	2
2.1	Desk Study	2
2.2	Survey Area	3
2.3	Survey guidance	3
2.4	Survey Dates	4
2.5	UK Habitat Classification, INNS and Hedgerows	4
2.6	Badger	5
2.7	Bats	6
2.8	Great Crested Newt	8
2.9	Limitations and Caveats	9
3.	Results	10
3.1	Desk Study	10
3.2	UK Habitat Classification, Hedgerows, and INNS Survey	12
3.3	Badger Survey	15
3.4	Bat Survey	17
3.5	Great Crested Newt Survey	22
Refe	rences	24

- Annex 1: Figures
- Annex 2: Botanical Species Lists
- Annex 3: Habitat Condition Assessment
- Annex 4: Bat Survey Results
- Annex 5 Photos of Habitats and Ecological Features

Appendix 3: Biodiversity Baseline

1. Introduction

1.1 Purpose of this Document

- 1.1.1 This appendix sets out the biodiversity baseline relevant to the proposed GSP substation. This appendix sets out the desk study, field survey methodology and results to provide a baseline of biodiversity information.
- 1.1.2 Specifically, it reports on:
 - UK Habitat Classification (UKHab) survey with habitat condition assessment;
 - Hedgerow survey;
 - Invasive Non-Native Species (INNS);
 - Bat surveys comprising preliminary ground level roost assessment (PGLRA) and subsequent aerial inspection through tree climbing and dusk emergence/dawn re-entry surveys;
 - Great crested newt (GCN) (*Triturus cristatus*) Habitat Suitability Index (HSI) assessment and environmental DNA (eDNA) surveys; and
 - Badger survey.
- 1.1.3 This appendix contained only baseline data. Any assessment of potential impacts on protected, priority and notable habitats and species, and recommendations on how these may be managed are discussed in the main Environmental Report. The appendix is supported by:
 - Annex 1 Figures, which illustrates results of the desk study and site surveys;
 - Annex 2 Plant Species and Abundance;
 - Annex 3 Habitat Condition Assessments;
 - Annex 4 Bat Survey Results; and
 - Annex 5 Photos of Habitats and Ecological Features.

1.2 Legislation

- 1.2.1 Legislation applicable to biodiversity is broadly split into two key types:
 - strict protection of sites or species; and
 - duties on all public bodies including local authorities, National Park Authorities and Natural England.
- 1.2.2 UK and English legislation relating to biodiversity, specific to the proposed GSP substation, includes:
 - The Environment Act 2021;
 - The Conservation of Habitats and Species Regulations 2017 (as amended);
 - The Wildlife and Countryside Act 1981 (as amended);
 - The Hedgerow Regulations 1997; and

• Natural Environment and Rural Communities (NERC) Act (2006).

2. Methodology

2.1 Desk Study

- 2.1.1 A desk study was undertaken to gather ecological information from publicly available and third-party sources. Information was gathered about the following ecological features:
 - Statutory and non-statutory designated sites (site plus 30km to identify Special Areas
 of Conservation (SAC) where bats are a qualifying feature and 2km for all other
 European sites, Sites of Special Scientific Interest (SSSI), National Nature Reserves
 (NNR), Local Nature Reserves (LNR) and locally designated sites or where there is a
 hydrological connection between the site and a designated site);
 - Protected and notable species (site plus 1km);
 - Habitats of principal importance i.e. 'priority habitats' and ancient woodland (site plus 1km); and
 - Invasive non-native species (site plus 1km).
- 2.1.2 Sources of information used were:
 - Descriptions of statutory designated sites were obtained from the Natural England website (Natural England, 2021);
 - Records of protected and notable species, including non-native invasive plant species were collated from the local Biological Records Centres (Suffolk Biodiversity Information Service (SBIS), Essex Wildlife Trust Biological Record Centre (EWTBRC), North East Essex Badger Group, and The Essex Field Club). SBIS and EWTBRC also provided descriptions and citations for non-statutory designated sites (Local Wildlife Sites (LoWS) in Essex);
 - Ecological surveys undertaken by The Environment Partnership (TEP) and Suffolk Wildlife Trust between 2010 and 2012 for the wider reinforcement project prior to project pause (unpublished), comprising habitats, bats, badger, breeding birds, hedgerow, GCN, and dormouse;
 - Historic records of ancient, veteran and notable trees were collated from the Woodland Trust's Ancient Tree Inventory (Woodland Trust, 2021);
 - The Forestry Commission's National Forest Inventory provided accurate boundaries for woodland habitats;
 - The Multi-Agency Geographic Information for the Countryside (MAGIC) website was reviewed to identify the locations of any statutory designated sites, Natural England's Ancient Woodland Inventory sites and Natural England's Priority Habitats Inventory sites – Habitats of Principal Importance;
 - MAGIC was also used to search for European Protected Species Licences, Natural England pond surveys results (2017-2019) for District Level Licencing (DLL) and GCN Risk Zones relating to GCN occurrence and the level of impact development is likely to have on this species;
 - Ordnance Survey data layers were used to provide accurate mapping of roads, rivers and waterbodies; and

• Pre-existing aerial imagery was assessed from the ESRI World Imagery dataset.

2.2 Survey Area

- 2.2.1 The proposed GSP substation is located in a predominantly rural landscape comprising arable farmland, surrounded by parcels of ancient woodland, the A131 and small rural roads.
- 2.2.2 The survey area is shown on Figure A3.1 in Annex 1 of this Appendix. To support the baseline information that will be submitted to support an application pursuant to Section 37 of the Electricity Act 1989 to the Department for Business, Energy and Industrial Strategy (BEIS) for some of the accompanying works described in Section 2 of the Environmental Report, the survey area is based on the on the proposed GSP substation location (i.e. 'the site'), the 400kV temporary overhead line diversion, the underground cables and the 132kV cable sealing end platform pylon. An additional 50m to 250m buffer was applied to the site and these accompanying works, depending on the biodiversity feature. The survey areas for the various types of biodiversity survey are detailed in Table 2.1.

Table 2.1: Defined survey areas for biodiversity field survey undertaken

Biodiversity Field Survey	Defined Survey Area
UKHab classification, INNS	The site, the 400kV temporary overhead line diversion, the underground cables, the 132kV cable sealing end platform pylon and a 50m buffer.
'Important' Hedgerow Survey	Hedgerows directly affected by the site, the 400kV temporary overhead line diversion, the underground cables, the 132kV cable sealing end platform pylon.
GCN HSI and eDNA	Water bodies within 250m of the site, the 400kV temporary overhead line diversion, the underground cables or the 132kV cable sealing end platform pylon.
Badger survey	The site, the 400kV temporary overhead line diversion, the underground cables, the 132kV cable sealing end platform pylon and a 50m buffer.
Bat surveys	The site, the 400kV temporary overhead line diversion, the underground cables, the 132kV cable sealing end platform pylon and a 50m buffer.

2.3 Survey guidance

- 2.3.1 The following guidance has been considered:
 - Guidelines for Preliminary Ecological Appraisal (CIEEM, 2017);
 - The UK Habitat Classification User Manual Version 1.1 (Butcher et al., 2020);
 - Defra Biodiversity Metric 2.0; Technical Supplement, Part 1 (Crosher et al., 2019);
 - Defra Biodiversity Metric 3.0; Technical Supplement, Part 1 (Panks et al., 2021);
 - Hedgerow Survey Handbook (Defra, 2007);
 - Bat Surveys for Professional Ecologists: Good Practice Guidelines (Collins, 2016); and
 - Evaluation of the Suitability of Habitat for the Great Crested Newt (Oldham et al., 2000).

2.4 Survey Dates

2.4.1 Table 2.2 lists the dates that field surveys were undertaken in 2021 and 2022. Surveys were undertaken as and when land access allowed, and within the appropriate survey season for each species/species group.

Table 2.2: Field survey dates

Biodiversity Field Survey	Defined Survey Area
UKHab classification, INNS, hedgerows	07/06/2021 – 11/06/2021, 23/06/2021 – 25/06/2021, 11/10/2021 – 14/10/2021
GCN HSI and eDNA	07/06/2021 - 11/06/2021, 23/06/2021 - 25/06/2021, 29/06/2021 - 30/06/2021
Badger survey	10/06/2021 - 11/06/2021, 23/06/2021 - 25/06/2021
Bat Survey – PGLRA	07/06/2021 - 11/06/2021
Bat survey – summer tree climbing inspection	26/07/2021 - 30/07/2021, 05/08/2021 - 06/08/2021, 19/08/2021 - 20/08/2021, 26/08/2021 - 27/08/2021
Bat Survey – emergence/re- entry	02/08/2021 - 05/08/2021, 23/08/2021 - 26/08/2021, 20/09/2021 - 24/-09/2021, 27/08/2021 - 29/09/2021
Bat Survey – tree climbing inspection for hibernation roosts	17/01/2022 - 21/01/2022, 14/02/2022 – 18/02/2022

2.5 UK Habitat Classification, INNS and Hedgerows

UKHab and INNS

- 2.5.1 The UKHab field surveys were undertaken in accordance with The UK Habitat Classification User Manual Version 1.1 (Butcher *et al.*, 2020). The principal aim of UKHab is to provide a system for recording and classifying habitats. The system has a primary hierarchy of five nested levels of complexity, from major ecosystems at level 1, ecosystem type at level 2, broad habitats at level 3, habitats including priority habitats at level 4, down to habitats including Annex 1 habitats at level 5. The primary code classification can be supplemented by secondary codes to provide more information. Some secondary codes are mandatory in the classification of certain habitat types such as ponds and traditional orchards.
- 2.5.2 Dominant and notable plant species were recorded, and plant abundance was categorised using the DAFOR scale; Dominant (D), Abundant (A), Frequent (F), Occasional (O) and Rare (R). Botanical taxonomic nomenclature follows that of the New Flora of the British Isles; Fourth Edition (Stace, 2019). Surveying was at the fine scale minimum mapping unit (MMU) i.e. 25m² polygons and linear features greater than 1m in width. However, hedgerows and other linear features were maintained as lines with their dimensions recorded.
- 2.5.3 During the habitat survey, the presence and extent of any Invasive Non-Native Species (INNS) were also recorded.
- 2.5.4 The habitats were then assessed on their condition. Habitat condition is a score based on the quality of the habitat, judged against the perceived ecological optimum state for that particular habitat, and criteria set out in the Natural England Biodiversity Metric 3.0 Guidance (DEFRA, 2021).

2.5.5 It should be noted that areas of cropland, intensive orchards and domestic garden are not required to have a habitat condition assessment. Urban sealed land and buildings have also been excluded.

Hedgerows

- 2.5.6 Hedgerows form part of the UKHab methodology. However, three hedgerows were identified as requiring more in-depth survey as they had potential to meet the biodiversity criteria for 'Important' hedgerows under the Hedgerow Regulations 1997 and where there is a need for hedgerow to be removed/damaged. The survey involved recording key characteristics such as physical attributes, species diversity, gaps in canopy, management and associated features such as adjacent banks, walls and ditches. The survey recorded the following in 30m sections:
 - Woody plants;
 - Climbing plants;
 - Woodland herbs, including those listed on Schedule 2 of the Hedgerows Regulations 1997 and ancient woodland indicators (Rose, 1999); and
 - Invasive non-native plants, such as Himalayan balsam (Impatiens glandulifera).

2.6 Badger

- 2.6.1 The survey followed Natural England survey guidance (Natural England, 2011). Signs of badger activity such as the presence of setts, dung pits, latrines, snuffle holes, tracks, hairs, prints and scratch marks were recorded. Particular attention was paid to areas where the vegetation and/ or topography offered suitable sett building sites for badgers, for example sloping banks and areas with dense ground cover (e.g. hedgerows, scrub and woodland). All field signs were recorded and plotted on a collector map on an iPad and photographs were taken as a record of the site visit.
- 2.6.2 Any setts found were mapped and the number of individual entrances and tunnel direction was recorded. Setts were provisionally categorised according to Natural England (2011), using the following criteria:
 - Main Sett: usually appearing well-used, well established, and having a large number of holes with big active spoil heaps, often with piles of old bedding outside. Main setts tend to have well-worn pathways between the sett and foraging areas and between sett holes. They are generally considered to be breeding setts (i.e. where cubs are most likely to be born) and are often in use all year round. A social group of badgers would typically only have one main sett within their territory.
 - Annex Sett: always close to the main sett and are usually connected by one or more obvious well-worn pathways. They consist of several holes but are not necessarily in use the whole time, even if the main sett is very active. Should a second litter of cubs be born within the social group, they are likely to be raised within the annex sett.
 - Subsidiary Sett: often have very few holes, are usually at least 50m from the main sett and do not have an obvious pathway connecting them to another sett. Subsidiary setts are not continuously active.
 - Outlier Sett: usually comprise of one or two holes with very little spoil outside (thus indicating that the tunnel system underground is not extensive), have no obvious pathway connecting them with another sett and are used only sporadically.

- 2.6.3 An indication of the level of activity at each sett was made as follows (Natural England, 2009):
 - Active:
 - In current use: Well-used sett entrances contain no debris or vegetation, are obviously regularly used and often show signs of having been recently excavated. Badger field signs such as latrines, guard hairs, dung pits or footprints are present.
 - Partially used: Setts or holes with entrances not in regular use and which have debris (twigs, leaves) around the entrance. However, they could potentially be used regularly in the future with little clearance necessary.
 - Disused:
 - Setts or holes showing signs of not having been in use for a considerable period and would not be used again without extensive clearance by a badger. There would be no other badger field signs at disused setts.

2.7 Bats

2.7.1 Bat surveys were led by licenced bat ecologists in accordance with the Bat Conservation Trust Bat Survey Good Practice Guidelines 3rd edition (Collins, 2016).

Preliminary Ground Level Roost Assessment (PGLRA) of Trees

- 2.7.2 Trees within the survey area were assessed for their potential to support roosting bats. The trees were observed from ground level during the day using a high-powered torch and binoculars to inspect features from all aspects. Potential roosting opportunities looked for included features such as:
 - Cavities within trunk and branches;
 - Woodpecker holes;
 - Knot and callus roles;
 - Rot holes and deadwood;
 - Damaged limbs;
 - Splits and cracks;
 - Severe weather i.e. frost cracks; and
 - Lifted bark.
- 2.7.3 Evidence of roosting bats was also searched for including:
 - Bat droppings, feeding remains and corpses (with notes made on quantity, freshness and type);
 - Dark staining below an access point that may be caused by bat faeces or urine; and
 - Staining around a hole that may be caused by the natural oils in bat fur.
- 2.7.4 Each tree was classified with a potential to support roosting bats, either high, moderate or low, according to the highest suitability Potential Roosting Feature (PRF) identified during the inspection as per Table 2.3. Trees with moderate and high potential were subject to further bat surveys as detailed in the following sections. Trees with negligible roosting potential were not recorded.

Table 2.3: Bat roosting potential

Suitability	Roosting Habitat (adapted from Collins, 2016)
Negligible	Negligible habitat features on site likely to be used by roosting bats.
Low	A tree with one or more potential roosting sites that can be used by individual bats opportunistically. However, these potential roost sites do not provide enough space, shelter, protection, appropriate conditions and/or suitable surrounding habitat to be used on a regular basis or by larger numbers of bats (i.e. unlikely to be suitable for maternity or hibernation).
Moderate	A tree with one or more potential roost sites that could be used by bats due to their size, shelter, protection, condition and surrounding habitat, but unlikely to support a roost of high conservation status (i.e. maternity or hibernation roost).
High	A tree with one or more PRFs that are obviously suitable for use by larger numbers of bats on a more regular basis and potentially for longer periods of time due to their size, shelter, protection, condition and surrounding habitat.

Aerial Inspection and Tree Climbing

- 2.7.5 Following the initial PGLRA, trees that were assessed as requiring further inspection at height were climbed as an alternative to dusk emergence and dawn re-entry surveys. A team of two suitably qualified ecologists accessed PRFs at height using a two-rope system. Once in a safe position within the tree, PRFs were inspected further with the use of an endoscope and a torch in search for bats, or evidence of bats. Following more indepth inspection of PRFs, some trees were then reassessed for their potential to support roosting bats with some trees being downgraded or upgraded based on these inspections.
- 2.7.6 Following on from the initial aerial inspection, trees were then subject to numerous climbs across the active season for bats: moderate trees being climbed twice and high potential trees being climbed on three separate occasions.

Bat Emergence and Re-entry Surveys

2.7.7 Trees that were unsafe to climb or did not require tree climbing due to their potential being unlikely to benefit from aerial inspection, were subject to dusk emergence and dawn reentry surveys whereby an ecologist observed the tree from ground level looking for bats exiting or accessing the PRFs at dusk or dawn, respectively. Bat calls were recorded on different bat detectors (Elekon Batlogger, Anabat Swift and Anabat Walkabouts) through the season. Infrared cameras were occasionally used to aid the bat survey and were used as a substitute surveyor. Where a roost was recorded during a dusk or dawn survey, the bat calls, and video footage where applicable, were analysed to confirm the bat species recorded in the field.

Hibernation Surveys

2.7.8 Trees that were assigned as having potential to support hibernating bats following the PGLRA and aerial inspections were subject to a further two climbing inspections: one visit in January 2022 and one visit in February 2022.

2.8 Great Crested Newt

2.8.1 Eleven waterbodies were identified within 250m the survey area with the potential to support GCN. Where access was permitted, waterbodies were subject to a Habitat Suitability Index (HSI) assessment to evaluate the suitability of each waterbody with regards to GCN and subsequently sampled for environmental DNA (eDNA) to assess presence/likely absence of GCN.

GCN Habitat Suitability Index

- 2.8.2 This methodology considers ten suitability indices that are known to affect GCN and their utilization of certain habitats. These indices comprise: geographical location; waterbody area; waterbody permanence; water quality; shade cover; waterfowl presence; fish presence; pond density/number of ponds within 1km; terrestrial habitat quality; and waterbody macrophyte cover.
- 2.8.3 Each waterbody was assessed and assigned a score between 0 and 1, with values closer to 0 being less suitable for GCN and those closer to 1 being more suitable. These values were then multiplied together, and the tenth root calculated to give an overall HSI score for the waterbody (Oldham *et al.*, 2000). Waterbodies were then allocated an associated predicted presence class, as shown in Table 2.4.

HSI score	Predicted presence class
0.00 - 0.49	Poor
0.50 – 0.59	Below average
0.60 - 0.69	Average
0.70 – 0.79	Good
0.80 – 1.00	Excellent

Table 2.4: Habitat Suitability Index assessment scores

GCN eDNA Sampling

2.8.4 Each waterbody was sampled for the presence/likely absence of GCN by collecting water samples which were subsequently sent to a laboratory for testing in the hope of identifying GCN DNA. Twenty water samples were taken from around the perimeter of each waterbody using a 40ml ladle, focusing on areas most likely to be used by GCN. The water samples were then transferred into a single Whirl-Pak bag. Before each sample was taken, the water was gently stirred using the ladle. This is because eDNA will often be present in larger quantities at the bottom of the waterbody as it tends to sink in water. The Whirl-Pak bag was then gently shaken to mix eDNA across the whole water sample. A pipette was then used to transfer 15ml of water from the Whirl-Pak bag into each of six conical tubes containing a preserving fluid. Each conical tube was then vigorously shaken for ten seconds to mix the water sample and the preservative. The six conical tubes were then NatureMetrics lab for analysis.

2.9 Limitations and Caveats

- 2.9.1 An absence of desk study records does not mean an absence of such habitats and species in that area. It may be due to under-recording. The desk study is designed to give an overview of the habitats and species previously recorded in the local area and provides an indication of which habitats and species may be present.
- 2.9.2 This report outlines the results of a snapshot in time, protected species such as badgers, bats and GCN are mobile and may utilise, or move into and out of, an area at short notice and as such may have been missed during the surveys or may be present at different times of the year. Additionally, populations of plants are often transient in nature and a survey undertaken over a narrow time frame can only provide an indication of the species present.
- 2.9.3 Some habitats were very densely vegetated, including dense bramble scrub in woodland areas that made them inaccessible to surveyors. However, the areas that could not be accessed were relatively limited in size and could be mostly viewed from their edges, making it unlikely that any features of ecological importance were missed.
- 2.9.4 Land access was not granted by landowners to certain areas, meaning these areas were unable to be surveyed. This included one pond (P1 on Figure A3.7 in Annex 1 of this Appendix)) that was not able to be assessed for HSI or sampled for eDNA).
- 2.9.5 Between the commencement of the first UKHab survey in June 2021 and the final habitat surveys in October 2021, the Natural England Biodiversity Metric was updated from v2.0 to v3.0 with revised guidance and criteria for habitat condition assessment. Following the release of the new version, the original habitat conditions assessed using version 2.0 were updated in line with version 3.0 criteria.
- 2.9.6 Hedgerow h2a-12 was not surveyed in the field. Therefore, based on aerial mapping and other hedgerows in the vicinity, and as a precaution, it has been classified as priority hedgerow in good condition. Additionally, based on a desk top study, it has been assessed as not meeting 'important' hedgerow status.

3. Results

3.1 Desk Study

Designated sites

- 3.1.1 There are no SACs within 30km of the site for which bats are a qualifying feature. There are no SACs, SPAs or Ramsar sites within 2km of the site or have any hydrological connection with the site. There are no NNR, SSSI or LNR within 2km of the site.
- 3.1.2 There are nine non-statutory locally sites located within 2km of the site, the closest being located directly adjacent to the proposed GSP substation. These are Butler's Wood LoWS and Waldegrave Wood LoWS, both of which are also ancient woodlands. Table 3.1 details the LoWS within 2km of the site. Figure A3.1 in Annex 1 shows the location of these sites.
- 3.1.3 The Ancient Woodland Inventory identified a number of sites of Ancient and Semi-Natural Woodland which were concurrent with LoWS designations. These are also detailed in Table 3.1.

LoWS name	Value	Features (adapted from EWT citations)	Approximate distance from site
Butler's Wood (Bra201)	National (included on the Ancient Woodland Inventory [AWI])	Similar in structure to Waldegrave Wood. An ancient woodland dominated by pedunculate oak <i>Quercus robur</i> . The site holds a wide variety of floral ancient woodland indicator species including the native bluebell <i>Hyacinthoides non-scripta</i> , yellow pimpernel <i>Lysimachia nemorum</i> , primrose <i>Primula vulgaris</i> , and wood anemone <i>nemorosa</i> .	Immediately north
Waldegrave Wood (Bra200)	National (AWI)	Similar in structure to Butler's Wood. An ancient woodland dominated by pedunculate oak and ash <i>Fraxinus excelsior</i> . With an understory containing hazel <i>Corylus avellana</i> coppice and containing several ancient woodland indicator species including the native bluebell, wood anemone, primrose and wood sorrel <i>Oxalis acetosellla</i> .	Immediately south
Twinstead Green (Bra210)	County	Damp grassland with a small pond and scattered trees, the site is noted to contain a moderately diverse assemblage of herb species.	600m ESE
Almshouse Wood (Bra211)	County	A woodland dominated by pedunculate oak, ash, birch, and hazel coppice. The ground flora has become dominated by bracken <i>Pteridium aquilinum</i> and bramble <i>Rubus fruticosus</i> <i>agg,</i> but ancient woodland indicators of primrose and dog's mercury <i>Mercurialis perennis</i> can be found.	670 NE
Parsonage Wood (Bra196	National (AWI)	An ancient woodland dominated by a birch <i>Betula sp</i> canopy and hazel understory, although there has been some replanting of conifer species historically. The site supports a wide variety of flora and bird species.	900m N
Twinstead Marsh (Bra222)	County	This site holds a wide variety of habitats including alder <i>Alnus glutinosa</i> and willow <i>Salix sp</i> carr, marsh and open water. These provide a variety of habitats for notable species such as nightingales <i>Luscinia megarhynchos</i> and willow warblers <i>Phylloscopus trochilus</i> .	1.38km E

Table 3.1: Non-statutory, locally designated sites within 2km of the site

LoWS name	Value	Features (adapted from EWT citations)	Approximate distance from site
Fenn Farm Mosaic (Bra223)	County	The site holds habitats ranging from willow carr, alder woodland, and marsh. The site supports a wide variety of wetland plant species.	1.34km E
Twinsteadhall Wood (Bra219	National (AWI)	Ancient woodland which has been replanted with coniferous species. The site holds a variety of ancient woodland ground flora species and supports a wide variety of invertebrate species.	1.5km SE
Coopersfield Wood (Bra228)	County	A small woodland dominated by pedunculate oak, ash, birch, and poplar <i>Populus sp.</i> The site supports a variety of invertebrate and bird species.	1.8km NE

3.1.4 Habitats of Principal Importance (HPI) identified from the Priority Habitat Inventory comprised two parcels of deciduous woodland associated with Bulter's and Waldegrave Woods. Beyond this and up to 1km, were additional HPIs comprising further parcels of deciduous woodland and traditional orchard. These can be seen on Figure A3.1 in Annex 1.

Protected, Priority and Notable Species

- 3.1.5 A summary of species records provided by Essex Wildlife Trust, Essex Field Club and North East Essex Badger Group is provided below.
- 3.1.6 A large number of bird records were returned within 1km from the site. These are summarised below and shown in Figure A3.2 of Annex 1, in accordance with the Birds of Conservation Concern (BoCC) list (version 5) (Stanbury et al., 2021). This details the population status of birds in the UK; accordingly, each species is categorised onto the Green, Amber or Red List, indicating an increasing level of conservation concern. Particular species highlights were: cuckoo (*Cuculus* canorus); fieldfare (*Turdus pilaris*) (also Schedule 1 species), greenfinch (*Chloris chloris*), house sparrow (*Passer domesticus*); lapwing (*Vanellus vanellus*); lesser redpoll (*Acanthis cabaret*), linnet (*Linaria cannabina*); merlin (*Falco columbarius*) (Also Schedule 1); mistle thrush (*Turdus viscivorus*); skylark (*Alauda arvensis*); starling (*Sturnus vulgaris*); woodcock (*Scolopax rusticola*); and yellowhammer (*Emberiza citronella*).
- 3.1.7 The scrub and hedgerow habitats along field margins and the A131 afford suitable conditions for priority species such as yellowhammer (Red List), linnet (Red List), house sparrow (Red List), starling (Red List), and mistle thrush (Red List).
- 3.1.8 Figure A3.3 in Annex 1 shows the spatial distribution of all other species records. Protected, priority and notable species records include brown long-eared bat (*Plecotus auritus*), common pipistrelle (*Pipistrellus pipistrellus*); badger (*Meles meles*) brown hare (*Lepus europaeus*) and great crested newt (*Triturus cristatus*).
- 3.1.9 A desk study search for European Protected Species (EPS) mitigation licences up to 2km from the site identified two bat licences, see Table 3.2.

Table 3.2: EPS Licence within 2km of the site

Species	Date	Case reference for Granted Application	Grid reference	Approximate distance to the site
Brown long-eared and common pipistrelle	2017-2027	2017-31391-EPS-MIT	TL862365	1.6km SE
Brown long-eared, common pipistrelle, soprano pipistrelle	2018-2024	2018-37898-EPS-MIT	TL86013850	1.8km NE

- 3.1.10 A desk study search for GCN records, using national Natural England pond surveys for District Level Licencing results between 2017-2019 and GCN Risk Zones, failed to identify any locations of GCN presence within 2km of the site. The site also falls within amber and green GCN risk zones. However, previous unpublished survey work by National Grid in 2012 at this location confirmed GCN presence within 250m of the site.
- 3.1.11 Dormouse surveys were undertaken by Suffolk Wildlife Trust in 2012 for the wider reinforcement project prior to project pause. The surveys included Butler's Wood, Waldegrave Wood and the hedgerow that connects the two areas of woodland adjacent to the A131. No dormice were recorded within these areas at that time. The nearest positive survey result was located 2.9km south east of the site. A precautionary approach is therefore being taken and dormice are assumed to be present where habitat is suitable to support them.

INNS

3.1.12 No records of INNS were returned within 1km of the site.

3.2 UK Habitat Classification, Hedgerows, and INNS Survey UKHab

3.2.1 The habitats recorded within the survey area during field survey are described below, grouped by broad habitat types, and Habitats of Principal Importance (HPI) identified. The location and distribution of these habitats (with their habitat parcel references) can be seen in Figure A3.4 in Annex 1. Table 3.3 summarises the habitats recorded and their condition assessment. Botanical species lists are provided in Annex B and a full breakdown of condition assessments is provided in Annex C of this Appendix.

Cropland (c)

3.2.2 The survey area was predominantly formed from c1 cropland (arable).

Hedgerow (h)

3.2.3 Hedgerows recorded within the survey area were primarily bordering arable land or adjacent to roads/access tracks. Seven hedgerows were recorded, all qualifying as priority hedgerows (habitat parcel references h2a-2, h2a-4, h2a-5 to ha-7, h2a-10 and h2a12).

Grassland (g)

3.2.4 The grassland areas recorded within the survey area were primarily other neutral grassland types along arable field margins (habitat parcel references g3c-6 and g3c-9). None of these grassland habitats are HPI.

Urban (u)

3.2.5 Urban habitat types recorded were roads - built linear features (u1e).

Woodland and Forest (w)

- 3.2.6 Woodland areas recorded within the survey area were primarily small areas of other broadleaved woodland types (habitat parcel references w1g7-1 to w1g7-5) but inclusive of the following HPI: Lowland Mixed Deciduous Woodland habitat habitat parcel reference: w1f7-1, w1f7-2.
- 3.2.7 One line of trees was recorded within the survey area (habitat parcel reference w1g6-2).

Habitat	Habitat Codes Figure A3.6	Condition	Notes
Lowland mixed deciduous woodland	w1f7-1 to w1f7-2	Good	Priority habitat and two parcels of ancient woodland – numerous ancient woodland indicator species recorded
Other broadleaved woodland	w1g7-1	Moderate	Small woodland and pond
Line of trees	w1g6-2	Moderate	Mature oak with some gaps and some semi- mature trees included
Other neutral grassland	g3c-9	Moderate	Arable field margin
	g3c-6	Poor	Road verges along A131, inclusive of adjacent nettle dominated boundary habitats
Hedgerow (priority)	h2a-2, h2a-4 to h2a- 7, h2a-10, h2a-12	Moderate to Good	Mostly agricultural hedges separating arable field, and alongside tracks. Well maintained with diverse species and adjacent ground flora intact

Table 3.3: UKHab parcels and conditions

Rivers and Lakes (r)

- 3.2.8 Within the survey area, no watercourses (rivers, streams, or lakes) were present. Ditches were present (see Annex 1 Figure A3.4) and surveyed (see Table 3.4) but they were dry and unlikely to hold water for more than four months of the year. Therefore, the ditches recorded did not meet the criteria of relevant primary level UKHab classification and have been classified only through secondary habitat types dry (code: 117) and ditch (code 191) shown as 'Dry ditch' on Figure A3.4 in Annex 1. Due to the ditches not holding water, all fail numerous criteria of the condition assessment automatically being assigned poor condition.
- 3.2.9 One pond is located within the 50m survey area, r1a6-3, but it does not qualify as a HPI. The pond was located within woodland with little submerged or riparian vegetation, was highly shaded and contained turbid water. As such it was assessed as being of poor condition.

Table 3.4: Summary of ditches recorded

Approximate Length (m)	Notes
130	Dry ditch, dominated by grasses, shallow gradient adjacent to road and overlapping with hedgerow H3 for short section.
270	Dry ditch, dominated by grasses. Some rush species. Willow and oak saplings. One mature oak.
1265	Dry ditch network, agricultural, approx. 1-1.5m wide, may fill with water at other times of year but not for over 4 months - substrate under dense bramble moist in some places, dominated by grasses with cow parsley, red campion, nettles present. Steep gradient.
330	Dry ditch running along line of trees, where one line stops the ditch continues unshaded and deeper - appeared artificially dug, and re-joined second line of trees along southern field boundary.
	Approximate Length (m) 130 270 1265 330

INNS

3.2.10 Whilst carrying out the UKHab survey, three areas of invasive non-native variegated archangel (*Lamium galeobdolon* subsp. *Argentatum*) were recorded, see Figure A3.4 in Annex 1 and photographs 44 and 45 in Annex 5.

Hedgerows

3.2.11 Full details of the condition of hedgerows within the 50m survey area can be seen in Annex 3. A summary of the hedgerow assessment is set out in Table 3.5. None of these hedgerows were classified as 'important' for biodiversity reasons under criteria 4 of the Hedgerow Regulations 1997.

Table 3.5: Hedgerow assessment summary

Hedgerow ID	Central grid reference	UKHab code & condition assessment	Hedgerow type	'Important' hedgerow under regulations Y/N	Justification
h2a-2	TL 84648 37136	h2a – good condition	Shrubby hedgerow	Ν	None of the criteria met – six woody species present but none of required associated features present
h2a-5	TL 84387 36700	h2a - good condition	Shrubby hedgerow	Ν	None of the criteria met – five woody species present but only one required associated feature present (parallel hedge within 15m)
h2a-6	TL 84392 36689	h2a - good condition	Shrubby hedgerow	Ν	None of the criteria met – five woody species present but only one required associated feature present (parallel hedge within 15m)

3.3 Badger Survey

3.3.1 Table 3.6 details the findings of the badger survey. Badger sett locations, mammal burrows, and badger field signs are shown in Annex 1 - Figure A3.5.

ID (see Figure A3.7)	Grid reference	Sign of current use - Y/N	Description	Likely sett type
B1	TL 84220 37204	Ν	Disused single entrance. Tunnel heading NW into bank, tunnel SE facing. D shaped hole with soil and debris making entrance shallow	Outlier
B2	TL 84241 37229	Y	One hole - spoil in front worn down by recent rain, looks occasionally used. Some potential scratch/slide marks into entrance. Guard hairs around entrance. Another disused hole nearby and well-worn path. Tunnel heading NE entrance SW facing	Outlier
B3	TL 84378 37160	Ν	Mammal burrow around fallen tree root system - rabbit dropping nearby	N/A
B4	TL 84448 37156	Ν	Two holes. Potential disused badger sett / active fox den, fox poo nearby. May have at one point been used by badger but not in current use. Some smaller rabbit burrows in vicinity.	Outlier
B5	TL 84465 37154	Ν	One potential badger entrance, no signs of current use. Tunnel on bank going NW direction facing SE	Outlier
B6	TL 84473 37162	Y	Badger entrance with hairs, spoil with some bedding in spoil, tunnel going N toward woodland - S facing, rabbit burrows nearby but only one showing signs of current use by badger	Outlier
B7	TL 84509 37167	Y	Badger sett two entrances - 1 around base and roots of aspen tree, tunnel goes N, entrance S facing, second entrance 2m west, tunnel goes NE, entrance SW facing -could be chamber in between. Hairs found around entrances and some recent spoil	Annex/ Subsidiary
B8	TL 84594 37197	Y	Partially collapsed sett (could be old rabbit warren) two main entrances, badger hairs outside and flattened spoil by rain. Numerous other collapsed holes/ tunnels surrounding	Annex/ Subsidiary
B9	TL 85165 37221	Y	Three sett entrances on bank, one showing large fresh spoil pile - two filled with loose debris. Clear pathway from track across ditch to holes. NW facing holes	Annex/ Subsidiary
B10	TL 84996 37124	Y	Badger sett under piles of bonfire wood. Two holes partially collapsed/ filled with debris from wood pile, one clear with some spoil. SE facing. Anecdotal evidence from member of public claimed wood recently dumped over existing sett so badgers dug new exits	Annex/ Subsidiary

Table 3.6: Badger survey results

ID (see Figure A3.7)	Grid reference	Sign of current use - Y/N	Description	Likely sett type
B11	TL 84905 37042	Y	Badger entrance under tree roots, very smoothed down, compacted soil, some old spoil with numerous badger hairs in	Outlier
B12	TL 84921 36958	Y	Badger entrance on bank. Active with fresh spoil heap, footprint, clear open tunnel headed SE, entrance W facing	Outlier
B13	TL 84913 36945	Y	Badger sett with recent spoil heaps, two holes either side of line of trees - E facing hole heading NW direction, hole adjacent to crop also E facing heading W, badger hair in spoil	Outlier
B14	TL 84953 36876	Y	Large main sett 23+ holes, 18 showing signs of current use - fresh spoil, recent footprint, hairs in spoil and surround, other entrances partially collapsed. 10 holes are within hedge likely more along length but obscured by dense vegetation	Main
B15	TL 84988 36907	Y	Single badger entrance 40m north west of main sett in field – likely connected as tunnel looks to head toward B14.	Annex/ Subsidiary Linked to B14
B16	TL 85015 36873	Ν	One disused badger entrance next to hedgerow, old spoil heap compacted with some vegetation growing through, partial debris in tunnel - hair found within compacted spoil	Outlier
B17	TL 85053 36871	Y	Badger entrance at base of mature oak tree trunk - hair and bark lifting found - entrance SW facing, and tunnel has NE direction to left of bark opening	Outlier
B18	TL 84092 36643	Ν	One potential badger entrance on side of road – no signs of activity	Outlier
B19	TL 84160 36533	Y	Main badger sett, at least 12 entrances in line of trees on bank with entrances in nearby grassland to east as well. Active signs fresh spoil and badger hairs, one latrine located nearby	Main
B20	TL 84146 36472	Y	Potential badger outlier fresh spoil outside	Outlier
B21	TL 84733 37118	Ν	Two holes one disused partially blocked, one with old spoil	Outlier

Incidental sightings

3.3.2 Numerous incidental sightings of live badger were recorded by surveyors carrying out bat dusk emergence and dawn re-entry surveys. The location of these can also be seen in Figure A3.5 in Annex 1.

3.4 Bat Survey

3.4.1 A summary of the bat surveys undertaken can be seen in Table 3.7 and full survey results of PGLRA is shown in Figure A3.6 of Annex 1. An analysis of bat call recordings is also given in Annex 4.

Preliminary Ground Level Bat Roost Assessment (PGLRA)

- 3.4.2 Forty four trees were recorded with potential to support roosting bats, see Figure A3.6 in Annex 1. No bats, or evidence of bat such as droppings, were recorded during the PGLRA.
- 3.4.3 Of these trees, 30 were identified as requiring further aerial inspection through tree climbing surveys. The remaining trees were identified as requiring dusk emergence and dawn re-entry surveys either because trees were unsafe to climb, or because the tree was unlikely to benefit from further inspection.

Bat Active Season Tree Climbing Surveys

- 3.4.4 During the tree climbing surveys, eleven trees had their level of bat roost potential either downgraded or upgraded depending on the internal characteristics of features recorded.
- 3.4.5 One confirmed roost was identified during the tree climbing surveys at TC27 (photo 32 Annex 5). At least two individuals of a *Myotis* bat species were identified through use of an endoscope camera. Bat droppings were collected and sent to a laboratory for analysis. The laboratory results confirmed that the droppings were from Natter's bat (*Myotis nattereri*).

Bat Emergence and Re-entry Surveys

- 3.4.6 One transitional roost of a soprano pipistrelle (*Pipistrellus pygmaeus*) was recorded in T16 (Photo 31 Annex 5). Please note, this tree is now located outside of the survey area. A single bat was recorded through infrared camera dropping down from a woodpecker hole on the northern aspect, flying around the tree and perching on the opposite side near an additional woodpecker hole. This footage was recorded at approximately 20:23 on 23 September 2021. Call sound analysis confirmed the species as being soprano pipistrelle, as seen in Annex 4.
- 3.4.7 No other bats were recorded emerging from, or re-entering, the trees during the dusk and dawn surveys.

Table 3.7: Bat survey results summary

Tree ID	Potential assigned at PGLRA	Potential following tree climbing surveys	Active Season Survey Type	Hibernation Potential Y/N	Hibernation surveys carried out Y/N	Roost recorded Y/N
TC1	Moderate	Moderate	Climbing inspection	Ν	N/A	Ν
TC2	Moderate	High	Climbing inspection	Y	Y	Ν
TC3	Moderate	Moderate	Climbing inspection	Y	N – Unsafe to climb	Ν
TC4	Moderate	Moderate	Climbing inspection	Ν	N/A	Ν
TC5	Moderate	Negligible	Climbing inspection- downgraded potential - no further survey	Ν	N/A	N/A
TC6	Moderate	Moderate	Climbing inspection	Ν	N/A	Ν
TC7	Moderate	Moderate	Climbing inspection	Ν	N/A	Ν
TC8	Moderate	Moderate	Climbing inspection	Y	Y	Ν
TC9	Moderate	Moderate	Climbing inspection	Ν	N/A	Ν
TC10	Moderate	Moderate	Climbing inspection	Y	Y	Ν
TC11	Moderate	Moderate	Climbing inspection	Ν	N/A	Ν
TC12	Moderate	Low	Climbing inspection- downgraded potential - no further survey	Ν	N/A	N/A
TC13	Moderate	Moderate	Climbing inspection	Ν	N/A	Ν
TC14	Moderate	Moderate	Climbing inspection	Y	Y	Ν
TC15	Moderate	High	Climbing inspection	Y	Y	Ν

Tree ID	Potential assigned at PGLRA	Potential following tree climbing surveys	Active Season Survey Type	Hibernation Potential Y/N	Hibernation surveys carried out Y/N	Roost recorded Y/N
TC16	Moderate	Moderate	Climbing inspection	Y	Y	Ν
TC17	Moderate	Low	Climbing inspection- downgraded potential - no further survey	Ν	N/A	N/A
TC19	Moderate	High	Climbing inspection	Y	Y	Ν
TC20	Moderate	Negligible	Climbing inspection- downgraded potential - no further survey	Ν	N/A	Ν
TC21	Moderate	Low	Climbing inspection- downgraded potential - no further survey	Ν	N/A	N/A
TC22	Moderate	Moderate	Climbing inspection	Y	Y	Ν
TC23	High	High	Climbing inspection	Y	Y – inspected from ground level	Ν
TC24	Moderate	Moderate	Climbing inspection	Ν	N/A	Ν
TC25	Moderate	High	Climbing inspection	Y	Y	Ν
TC26	Moderate	Low	Climbing inspection	Ν	N/A	N/A
TC27	Moderate	Moderate	Climbing inspection	Y	Y	Y
TC28	Moderate	High	Climbing inspection	Y	Y	Ν
TC29	Moderate	Moderate	Climbing inspection	Y	Y	Ν
TC32	N/A - not recorded at PGLRA	High	Climbing inspection	Y	Y	Ν
TC33	N/A - not recorded at PGLRA	Moderate	Climbing inspection	Ν	N/A	Ν

Tree ID	Potential assigned at PGLRA	Potential following tree climbing surveys	Active Season Survey Type	Hibernation Potential Y/N	Hibernation surveys carried out Y/N	Roost recorded Y/N
T1	High	N/A - climb not required	Emergence/re-entry survey	Y	Y – inspected from ground level	Ν
T2	Moderate	N/A - climb not required	Emergence/re-entry survey	Y	Y – no roost recorded	Ν
Т3	Moderate	N/A - climb not required	Emergence/re-entry survey	Ν	N/A	Ν
Τ4	High	N/A - climb not required	Emergence/re-entry survey	Y	Y – inspected from ground level	Ν
T5	Moderate	N/A - climb not required	Emergence/re-entry survey	Ν	N/A	Ν
Т6	Moderate	N/A - climb not required	Emergence/re-entry survey	Y	Y – inspected from ground level	Ν
Τ7	High	N/A - climb not required	Emergence/re-entry survey	Y	N – unsafe to climb	Ν
Т8	High	N/A - climb not required	Emergence/re-entry survey	Y	N – unsafe to climb	Ν
Т9	High	N/A - climb not required	Emergence/re-entry survey	Υ	Y – inspected from ground level	Ν
T10	Moderate	N/A - climb not required	Emergence/re-entry survey	Y	N – unsafe to climb – bark checked from ground level where possible	Ν
T11	Moderate	N/A - climb not required	Emergence/re-entry survey	Ν	N/A	Ν
T12	Moderate	N/A - climb not required	Emergence/re-entry survey	Ν	N/A	Ν

Tree ID	Potential assigned at PGLRA	Potential following tree climbing surveys	Active Season Survey Type	Hibernation Potential Y/N	Hibernation surveys carried out Y/N	Roost recorded Y/N
T13	Moderate	N/A - climb not required	Emergence/re-entry survey	Ν	N/A	Ν
T15	Moderate	N/A - climb not required	Emergence/re-entry survey	Ν	N/A	Ν

- 3.4.8 Bat activity within the survey area was recorded during the emergence and re-entry surveys, with the most activity being recorded within Butler's Wood (habitat parcel reference: w1f7-1) and Waldegrave Wood (habitat parcel ref: w1f7-2).
- 3.4.9 Species recorded foraging and commuting during the surveys included:
 - Common pipistrelle (Pipistrellus pipistrellus);
 - Soprano pipistrelle;
 - *Myotis* species;
 - European barbastelle (Barbastella barbastellus);
 - Noctule (Nyctalus noctula); and
 - Brown long eared bat (*Plecotus auritus*).
- 3.4.10 There was notable activity of barbastelle bat in Butler's Wood and Waldegrave Wood (habitat parcel references: w1f7-1 and w1f7-2 on Figure A3.4 of Annex 1). Barbastelle bat was recorded foraging and commuting in both woodlands across all three survey visits. On occasion during surveys between 3rd and 5th August surveyors were able to gain sight of three to four barbastelle bats foraging together in the canopy.

Hibernation Surveys

- 3.4.11 No bat roosts were recorded during the hibernation surveys. Some of the trees had become unsafe to climb and so could not be fully inspected for hibernating bats.
- 3.4.12 A summary of the bat roosts identified is given in Table 3.8. Photographs of the tree roosts are in Annex 5 (Photos 30-32).

Tree ID	Date	Roost type	Roost feature	Species	No. of bats
TC27 Photo 32 – Annex E	20/08/2021 – summer tree climb and inspection	Summer day roost	Knot hole on main stem facing west	Natterer's bat <i>Myotis nattereri</i>	2-3
*T16 Photo 31 – Annex E	23/09/2021 - dusk emergence	Transitional	Woodpecker hole on main stem facing west	Soprano pipistrelle <i>Pipistrellus pygmaeus</i>	1

Table 3.8: Bat roosts recorded

*Now outside of survey area

3.5 Great Crested Newt Survey

3.5.1 The results of the GCN eDNA and HSI surveys are summarised in Table 3.9 and shown in Figure A3.7 of Annex 1. Two waterbodies (P5 and P6) returned positive tests for GCN eDNA, confirming GCN presence. One waterbody (P11) returned inconclusive¹ results. Photos of waterbodies testing positive for GCN eDNA can be seen in Annex 5.

¹ An inconclusive result can be issued for numerous of reasons: DNA from the great crested newt has not been detected but the controls have indicated that the sample has been degraded; DNA from the great crested newt has not been detected but the controls have indicated that the PCR has been inhibited. Additionally an inconclusive result may be due to the water chemistry, sediment or other debris in the sample. Therefore inconclusive findings cannot be used as definitive result to cause either presence or absence.

3.5.2 One waterbody (P1) could not be surveyed due to land access restriction(see Section 2.9). Four ditches were found upon initial survey to be dry and therefore unsuitable for GCN.

Table 3.9: GCN HSI and eDNA results

Waterbody name	Grid reference	Description	HSI score	HSI class	eDNA result
P1	Access denied				
Ρ5	TL 84459 37332	Large waterbody. Leaf litter dense on shallow margins, little aquatic vegetation, maybe some signs of pollution from oil on surface, steeper banks on west side, bracken and common nettle on banks, ducks observed.	0.79	Good	Positive
P6	TL 84479 37254	L shaped waterbody, some duckweed cover and small rush, some deadwood present, shallow bank, bramble bracken nettle on banks, common blue damselfly observed.	0.78	Good	Positive
P7	TL 84733 37130	Waterbody in copse of willow and oak trees, very little submerged or marginal vegetation, fallen branches and deadwood in waterbody, shallow banks not very vegetated apart from common nettle.	0.75	Good	Negative
P8	TL 84898 37301	Dry ditch in arable field, has two 90-degree bends, steep grassy banks and occasional trees.	N/A	N/A	N/A
P9	TL 84881 37145	Small, isolated waterbody in arable field, lots of catkins and leaf litter from surrounding willow trees, and algae in water.	0.5	Below average	Negative
P10	TL 84520 36950	Dry L shaped ditch, steep banks, surrounded by ferns, likely seasonally filled with water for short time.	N/A	N/A	N/A
P11	TL 84331 36969	Waterbody in woodland, shallow banks, surrounded by silver birch, algal bloom and signs of pollution, no aquatic vegetation.	0.55	Below average	Inconclusive
P12	TL 84218 37180	Dry ditch bordering woodland and arable field, steep grass banks.	N/A	N/A	N/A
P13	TL 84076 36382	Long dry ditch running adjacent to arable field with steep banks and line of trees on one of the banks – hazel, oak and hawthorn.	N/A	N/A	N/A
P23	TL 84344 37285	Small woodland waterbody, shallow with shallow bank, little to no vegetation.	0.47	Poor	Negative

References

Altringham J. (2003) British Bats. London, UK.

Biggs, J., Edwald, N., Valentini, A., Gaboriaud, C.,Griffiths, R.A., Foster, J. Wilkinson, J., Arnett A, Williams, P., and Dunn, F. (2014). Analytical and Methodological Development for Improved Surveillance of the Great Crested Newt. Defra Project WC1067. Freshwater Habitats Trust: Oxford, UK.

Butcher, B., Carey, P., Edmonds, R. Norton, L. and Treeweek, J. (2020) The UK Habitat Classification User Manual 1.1 at http://www.ukhab.org/ [accessed June 2021]

Butcher, B., Carey, P., Edmonds, R. Norton, L. and Treeweek, J. (2020) The UK Habitat Classification Field Key at http://www.ukhab.org/ [accessed June 2021]

Chartered Institute of Ecology and Environmental Management (2017) Guidelines for Preliminary Ecological Appraisal. Revised 3rd Edition. CIEEM, Winchester.

Collins, J. (ed) (2016). Bat Surveys for Professional Ecologists: Good Practice Guidelines. (3rd Edition). Bat Conservation Trust, London, UK.

Clark (2007). Badgers. Whittet Books, Suffolk, UK.

Defra (2019) Biodiversity Metric 2.0 Calculation Tool (December 2019 update). Available at http://publications.naturalengland.org.uk/publication/5850908674228224 [accessed 06/06/2021]

Defra (2019) The Biodiversity Metric 2.0 – User Guide – Beta Test. Available at http://publications.naturalengland.org.uk/publication/5850908674228224 [accessed 06/06/2021]

Defra (2019) The Biodiversity Metric 2.0 – Technical Supplement – Beta Test. Available at http://publications.naturalengland.org.uk/publication/5850908674228224 [accessed 06/06/2021]

Defra (2019) Biodiversity Metric 3.0 Calculation Tool. Available at http://publications.naturalengland.org.uk/publication/6049804846366720 [accessed 15/09/2021]

Defra (2021) The Biodiversity Metric 3.0 – User Guide – Beta Test. Available at http://publications.naturalengland.org.uk/publication/6049804846366720 [accessed 15/09/2021]

Defra (2021) The Biodiversity Metric 3.0 – Technical Supplement . Available at http://publications.naturalengland.org.uk/publication/6049804846366720 [accessed 15/09/2021]

DEFRA(2007). Hedgerow Survey Handbook: a standard procedure for local surveys in the UK. London, UK.

MAGIC (2021) MAGIC on the MAP. Online at: https://magic.defra.gov.uk/magicmap.aspx [Accessed October 2021]

National Grid (2022) Bramford to Twinstead – 2.1 Preliminary Environmental Information Report. Issue number: BT-JAC-560-0007.

Natural England (2009). Guidance on 'Current Use' in the definition of a Badger sett. Natural England, UK.

Natural England (2011). Badgers and Development – A Guide to Best Practice and Licensing. Natural England Interim Document Revised 12/11. Natural England, UK.

Natural England (2021) Designated Sites View. Available at: https://designatedsites.naturalengland.org.uk [Accessed 10/10/21/]

Natural England (2021). Ancient Woodland (England). [Online] Available at: https://data.gov.uk/dataset/9461f463-c363-4309-ae77-fdcd7e9df7d3/ancient-woodland-england [Accessed 05/09/21]

Oldham, R.S., Keeble, J., Swan, M.J.S. and Jeffcote, M. (2000). Evaluation of the Suitability of Habitat for the Great Crested Newt (Triturus cristatus). Herpetological Journal 10 (4), 143-155.

Rose, D., O'Reilly C., Collings, c. (2006) The Wildflower Key. London UK

Russ, J. (2012). British Bat Calls: A Guide to Species Identification. Pelagic Publishing. Exeter, UK.

Stace, C. (2019). New Flora of the British Isles (4th Ed), Cambridge University Press. London, UK.

Stanbury, A., Eaton, M., Aebischer, N., Balmer, D., Brown, A., Douse, A., Lindley, P., McCulloch, N., Noble, D. and Win, I. (2021) The status of our bird populations: the fifth Birds of Conservation Concern in the United Kingdom, Channel Islands and Isle of Man and second IUCN Red List assessment of extinction risk for Great Britain. British Birds (114) 723-747

Woodland Trust (2021) Ancient Tree Inventory. Available at https://ati.woodlandtrust.org.uk [Accessed 10/10/21/]

Annex 1: Figures

Figure A3.1 – Designated Sites and Priority Habitats

- Figure A3.2 Desk Study Birds
- Figure A3.3 Desk Study Excluding Birds
- Figure A3.4 UK Habitat, INNS and Hedgerow Survey Results
- Figure A3.5 Badger Survey Results
- Figure A3.6 Bat Survey Results
- Figure A3.7 Great Crested Newt Survey Results



Path: \\gblon0vs01\GISProj\B2600017_NG_Bramford2Twinstead\GIS\04_Environmental\09_Substation_ProtectedSpecies\AAA_B2T_GSP_Protected_Species_v2.aprx



Path: \\gblon0vs01\GISProj\B2600017_NG_Bramford2Twinstead\GIS\04_Environmental\09_Substation_ProtectedSpecies\AAA_B2T_GSP_Protected_Species_v2.aprx



Path: \\gblon0vs01\GISProj\B2600017_NG_Bramford2Twinstead\GIS\04_Environmental\09_Substation_ProtectedSpecies\AAA_B2T_GSP_Protected_Species_v2.aprx



Path: \\gblon0vs01\GISProj\B2600017_NG_Bramford2Twinstead\GIS\04_Environmental\09_Substation_ProtectedSpecies\AAA_B2T_GSP_Protected_Species_v2.aprx

	FIGURE A3.4
Legend	
Site	Boundary
50m	n Survey Area
	S - Variegated Yellow Archangel
Habitat type ((UK Hab)
—— w1g	6 - Line of trees
Dry	Ditch
— — h2a	- Hedgerow (priority habitat)
<u> </u>	- Built linear features
primary	
c1-	arable and horticulture
g3c	- other neutral grassland
g4 -	modified grassland
r1a6	5 - Other eutrophic standing waters
u1e	- built linear features
w1fl (pric	7 - Other Lowland mixed deciduous woodland ority habitat
F	The set
The second	
,	and the second sec
	Sudbury
	Land I have a
1	
1	
~	Nº TOL
	R. A.
	A I Ken
-	Heltrad
~	Halstead
	the that
0 04/2022	Final JT FM LG
Rev. Date	Purpose of revision Drawn Check'd Rev'd A
	Jacobs
	160 Dundee St, Edinburgh, EH11 1DQ, UK. Tel: +44(Q)131 659 1500
Client	
	national arid
	3
Project	
	PROPOSED GSP SUBSTATION OFF THE A131
Drawing Title	
Uł	K Habitat, Invasive Non-Native Species and Hedgerows
	Page 1 of 1
Drawing Status	Final
Scale @ A3	1:5,000 DO NOT SCA
Jacobs No. Client No.	BT-NG-020627-560-0032
Drawing No.	AAA_B2T_GSP_Protected_Species_Fig3.4_Rev01
© Copyright 20 document are t without the writ	121 Jacobs U.K. Limited. The concepts and information contained in this he property of Jacobs. Use or copying of this document in whole or in part ten permission of Jacobs constitutes an infringement of copyright.
Limitation: This Client, and is su	drawing has been prepared on behalf of, and for the exclusive use of Jacobs' bject to, and issued in accordance with, the provisions of the contract betwee Client Jacobs accents no liability or responsibility whether or is present
any use of, or re	eliance upon, this drawing by any third party.

Figure A3.5 (Badger Survey Results) contains confidential information and has been provided to Braintree District Council only



Path: \\gblon0vs01\GISProj\B2600017_NG_Bramford2Twinstead\GIS\04_Environmental\09_Substation_ProtectedSpecies\AAA_B2T_GSP_Protected_Species_v2.aprx

Site Boundary Bat Potential Tree Confirmed Roost High Moderate Low Negligible 	
Site Boundary Bat Potential Tree Confirmed Roost High Moderate Low Negligible Negligible	
Bat Potential Tree	
 Confirmed Roost High Moderate Low Negligible 	
 ▲ High ▲ Moderate ▲ Low ▲ Negligible 	
 Moderate Low Negligible 	
 Low Negligible 	/
Megligible	
Sudbury Halstead	
Sudbury Halstead	1
Halstead	· · · ·
0 04/2022 Final JT FM L Rev. Date Purpose of revision Drawn Check'd Rev.	LG ev'd
Jacobs	
160 Dundes St, Edinburgh, EH11 1DQ, UK. Tal: +4//0131 559 1500	
client nationalgrid	
PROPOSED GSP SUBSTATION OFF THE A131	
Bat Survey Results Page 1 of 1	
Drawing Status Eincl	
Scale @ A3 1:5,000 DO NOT	T SCA
Jacobs No. B2416601 Client No. BT-NG-020627-560-0032	
Drawing No. AAA_B2T_GSP_Protected_Species_Fig3.6 Rev0	-



Path: \\gblon0vs01\GISProj\B2600017_NG_Bramford2Twinstead\GIS\04_Environmental\09_Substation_ProtectedSpecies\AAA_B2T_GSP_Protected_Species_v2.aprx
Annex 2: Botanical Species Lists

Table 1: Lowland mixed deciduous woodland - HPI – Butler's Wood Ancient Woodland - w1f7-1

Common name	Latin Name	Abundance
Canopy Layer		
Oak	Quercus robur	D
Hazel	Corylus avellana	D
Ash	Fraxinus excelsior	A
Silver Birch	Betula pendula	F
Field maple	Acer campestre	F
Holly	llex aquifolium	0
Wild cherry*	Prunus avium	0
Hawthorn	Crataegus monogyna	0
Goat willow	Salix caprea	0
Small leaved lime*	Tilia cordata	R
Dogwood	Cornus sanguinea	R
Elder	Sambucus nigra	R
Ground Flora		
Wood anemone*	Anemone nemororsa	A
Bluebell*	Hyacinthoides non-scripta	A
Bracken	Pteridium	A
Cleavers	Galium aparine	A
Nettles	Urtica dioica	A
Wood sorrel*	Oxalis acetosella	F
Wood sedge	Carex sylvatica	F
Garlic mustard	Alliaria petiolata	F
Doc sp.	Rumex sp.	F
Speedwell germander	Veronica chamaedrys	F
Bugle	Ajuga reptans	F
Bramble	Rubus fruticosus	F
Ground ivy	Glechoma hederacea	F
Tufted hair grass	Deschampsia cespitosa	F
Primrose*	Primula vulgaris	F
Red campion*	Silene dioica	F
Honeysuckle	Lonicera periclymenum	F
Creeping bent	Agrostis stolonifera	F

Herb Robert	Geranium robertianum	F
False oat-grass	Arrhenatherum elatius	0
Yellow pimpernel*	Lysimachia nemorum	0
Thistle	Cirsium vulgare	0
Greater stitchwort	Stellaria holostea	0
Heath speedwell	Veronica officinalis	0
Fern	Tracheophyta	0
Pendulous sedge*	Carex pendula	0
Black bryony	Dioscorea communis	R
Pignut*	Conopodium majus	R
Soft brome	Bromus hordeaceus	R
Cocks foot	Dactylis glomerata	R

* ancient woodland indicators as per The Wildflower Key (Rose et al., 2006) for east England

Table 2: Lowland mixed deciduous woodland – HPI - Waldegrave Wood Ancient Woodland - w1f7-2

Common name	Latin Name	Abundance
Canopy layer		
Oak	Quercus robur	A
Hazel	Corylus avellana	F
Silver Birch	Betula pendula	F
Ash	Fraxinus excelsior	F
Aspen	Populus tremuloides	F
Blackthorn	Prunus spinosa	0
Wild cherry*	Prunus avium	0
Field maple	Acer campestre	R
Ground flora		
Male fern*	Dryopteris filix-mas	A
Bluebell*	Hyacinthoides non-scripta	A
Foxglove*	Digitalis purpurea	F
Honeysuckle	Lonicera periclymenum	F
Bracken	Pteridium aquilinum	F
Bramble	Rubus fruticosus	F
Cleavers	Galium aparine	F
Wood sorrel*	Oxalis acetosella	F
Wood anemone*	Anemone nemorosa	F

Speedwell germander	Veronica chamaedrys	F
Herb Robert	Geranium robertianum	F
Wood meadow grass*	Poa nemoralis	F
Rough meadow grass	Poa trivialis	F
Ground Ivy	Glechoma hederacea	F
Yellow pimpernel*	Lysimachia nemorum	0
Bugle	Ajuga reptans	0
Wood sedge*	Carex sylvatica	0
Pendulous sedge*	Carex pendula	0
Wood dock	Rumex sanguineus	0
Curled dock	Rumex crispus	0
Greater stitchwort	Stellaria holostea	0
Yorkshire fog	Holcus lantus	0
Creeping bent	Agrostis stolonifera	0
Garlic mustard	Alliaria petiolata	0
Tufted hair grass	Deschampsia cespitosa	R
Red campion*	Silene dioica	R
Black bryony	Dioscorea communis	R

*Shows ancient woodland indicators as per The Wildflower Key (Rose et al., 2006) for east England

Table 3: Other neutral grassland - parcel g3c-9 - Marginal grass between woodlands and arable

Common name	Latin Name	Abundance
False oat grass	Arrhenatherum elatius	A
Cocks foot	Dactylis glomerata	A
Common bent	Agrostis capillaris	A
Dandelion	Taraxacum officinale	F
Soft brome	Bromus hordeaceus	F
Thistle	Cirsium vulgare	F
Yorkshire Fog	Holcus lantus	F
Dock sp.	Rumex sp.	F
Germander speedwell	Veronica chamaedrys	0
Ragwort	Jacobaea vulgaris	0
Heath speedwell	Veronica officinalis	0
Creeping buttercup	Ranunculus repens	0

Perennial rye grass	Lolium perenne	0
Crested dog tail	Cynosurus cristatus	0
Birds foot trefoil	Lotus corniculatus	0
Common vetch	Vicia sativa	0
Goatsbeard	Aruncus dioicus	0

Table 4: Hedgerow (priority) - h2a-2 (H1) - Hedgerow between Butler's and Waldegrave woodlands

Common name	Latin Name	Abundance
Hawthorn	Crataegus monogyna	F
Blackthorn	Prunus spinosa	F
Field maple	Acer campestre	F
Hazel	Corylus avellana	F
Dog rose	Rosa canina	0
Oak	Quercus robur	0
Ground Flora		
False oat	Arrhenatherum elatius	F
Cocksfoot	Dactylis Glomerata	F
Ribwort plantain	Plantago lanceolata	0
Barron brome	Bromus sterilis	0
Yarrow	Achillea millefolium	0
Daisy	Bellis perennis	0
Bristly oxtongue	Helminthotheca echioides	R
Hedgerow Cranesbill	Geranium pyrenaicum	R

Table 5: Dry ditch - D9 – Dry ditch running adjacent and linking Butler's Wood and Waldegrave wood

Common name	Latin Name	Abundance
False oat grass	Arrhenatherum elatius	F
Bramble	Rubus fruticosus	F
Cow parsley	Anthriscus sylvestris	F
Creeping Buttercup	Ranunculus repens	F
Cocks foot	Dactylis glomerata	F

Broom	Sarothamnus scoparius	0
Rosebay willow herb	Chamaenerion angustifolium	0
Red campion	Silene dioica	0

Table 6: Line of trees – w1g6-2 - Along ditch D12 and part of Old Road

Common name	Latin Name	Abundance
Oak	Quercus robur	F
Ash	Fraxinus excelsior	F
Field maple	Acer campestre	F
Blackthorn	Prunus spinosa	0
Grey Willow	Salix cinerea	0

Table 7: Hedgerows	(priority)	- h2a-5 (H	2) and h2a-6	(H3) - north	h & south of Old	Road
--------------------	------------	------------	--------------	--------------	------------------	------

Common name	Latin Name	Abundance
Canopy layer		
Hawthorn	Crataegus monogyna	A
Field maple	Acer campestre	A
Hazel	Corylus avellana	A
Elder	Sambucus nigra	F
lvy	Hedera helix	F
Dogwood	Cornus sanguinea	0
Ground flora		
Cow parsley	Anthriscus sylvestris	F
Cocks foot	Dactylis glomerata	F
Common bent	Agrostis capillaris	F
False oat-grass	Arrhenatherum elatius	F
Creeping buttercup	Ranunculus repens	F
Nettles	Urtica dioica	0
Small flowered cranesbill	Geranium pusillum	0
Soft brome	Bromus hordeaceus	0
Germander speedwell	Veronica chamaedrys	0
Ragwort	Jacobaea vulgaris	0
Ribwort plantain	Plantago lanceolata	0
Greater stitchwort	Stellaria holostea	R

Table 8: Other broadleaved woodland - w1g7-1 – surrounding pond r1a6-2

Common name	Latin Name	Abundance
Canopy layer		
Crack willow	Salix fragilis	F
Hazel	Corylus avellana	F
Field maple	Acer campestre	F
Oak	Quercus robur	0
Hawthorn	Crataegus monogyna	0
Ground flora		
Nettles	Urtica dioica	A
Cleavers	Galium aparine	F

Meadow grass	Poa annua	0
Dog rose	Rosa canina	0

Annex 3: Habitat Condition Assessment

1. Grassland condition

Habitat Description - Grassland

Includes both agricultural, recreational, amenity, road verges and semi-natural grassland types including Priority Habitat Grasslands on all soil types.

Will be dominated by grassland species with very little (if any) dwarf shrub, wetland or wooded species within the sward.

Will exist above and below the level of enclosure at all altitudes

	Habitat Parce Figure A3.4)	I reference (see
Assessment Criteria	G3c-6	G3c-9
The appearance and composition of the vegetation closely matches characteristics of the specific grassland habitat type (see UKHab definition). Wildflowers, sedges and indicator species for the specific grassland habitat type are very clearly and easily visible throughout the sward.	✓	✓
Sward height is varied (at least 20% of the sward is less than 7 cm and at least 20 per cent is more than 7 cm) creating microclimates which provide opportunities for insects, birds and small mammals to live and breed.	Х	✓
Cover of bare ground between 1% and 5%, including localised areas, for example, rabbit warrens.	Х	\checkmark
Cover of bracken less than 20% and cover of scrub (including bramble) less than 5%.	✓	√
There is an absence of invasive non-native species (as listed on Schedule 9 of WCA, 1981). Combined cover of undesirable species1 and physical damage (such as excessive poaching, damage from machinery use or storage, damaging levels of access, or any other damaging management activities) accounts for less than 5% of total area.	x	Х

Habitat Quality	Assessment Criteria	Qualifying Habitat Parcels
Good	Passes 5 of 5 criteria	None
Moderate	Passes 3 or 4 of 5 criteria	G3c-9
Poor	Passes 0, 1 or 2 of 5 criteria	G3c-6

2. Woodland condition

Habitat Description - Woodland

Woodland is defined as vegetation dominated by trees more than 5 m high when mature, which forms a distinct, although sometimes open, canopy [areas of trees with a canopy greater than 20%]. This includes felled, young, or newly planted woodland.

There is no minimum size for areas of trees that have the definite characteristics and feel of a woodland and are managed as woodland.

Two broad woodland types are considered here:

- Broadleaved, mixed and yew woodland.
- Coniferous woodland.

It does not include scrub (see separate scrub condition assessment).

In England, native woodland is defined as woodland that is composed of at least 80% native tree species including 'naturalised species'. It is based on the England Woodland Biodiversity Group condition assessment for non SSSI woodlands.

Wood Pasture and Parkland

Wood pasture is a vegetation structure rather than a particular plant community. Typically, this structure consists of large, open-grown or high forest trees (often pollards) at various densities, in a matrix of grazed grassland, heathland and/or woodland floras.

This feature includes:

Wood pasture and parkland derived from medieval forests and embankments, wooded commons, parks and pastures with trees; and where the land use has been converted to arable, forestry or amenity, but where ancient trees are still present.

	Score		Habitat Parcel referenc (see Fig A3.4)		
Indicator	Good (3 points)	Moderate (2 points)	Poor (1 point)	W1f7- 1	W1f7- 2
Age distribution of trees	Three age classes present	Two age classes present	One age class present	3	3
Wild, domestic and feral herbivore damage	No significant browsing damage evident in woodland	Evidence of significant browsing pressure is present in 40% or less of whole woodland	Evidence of significant browsing pressure is present in 40% or more of whole woodland	3	3
Invasive plant species	No invasive species present in woodland	Rhododendron or laurel not present, other invasive species < 10% cover	Rhododendron or laurel present, or other invasive species > 10% cover	3	2

	Score			Habitat Parcel referen (see Fi A3.4)	t ice gure
Indicator	Good (3 points)	Moderate (2 points)	Poor (1 point)	W1f7- 1	W1f7- 2
Number of native tree species	Five or more native tree or shrub species found across woodland parcel	Three to four native tree or shrub species found across woodland parcel	None to two native tree or shrub species across woodland parcel	3	3
Cover of native tree and shrub species	> 80% of canopy trees and >80% of understory shrubs are native	50-80% of canopy trees and 50-80% of understory shrubs are native	< 50% of canopy trees and understory shrubs are native	3	3
Open space within woodland	10 – 20% of woodland has areas of temporary open space, unless woodland is 10ha in which case lower threshold of <10% does not apply	21- 40% of woodland has areas of temporary open space	More than 40% of woodland has areas of temporary open space	3	3
Woodland regeneration	All three classes present in woodland; trees 4- 7cm dbh, saplings and seedlings or advanced coppice regrowth	One or two classes only present in woodland	No classes or coppice regrowth present in woodland	3	3
Tree health	Tree mortality less than 10%, no pests or diseases and no crown dieback	11% to 25% mortality and/or crown dieback or low risk pest or disease present	Greater than 25% tree mortality and or any high risk pest or disease present	2	2
Vegetation and ground flora	Ancient woodland flora indicators present	Recognisable NVC plant community present	No recognisable NVC community	3	3
Woodland vertical structure	Three or more storeys across all survey plots or a complex woodland	Two storeys across all survey plots	One or less storey across all survey plots	2	2
Veteran trees	Two or more veteran trees per hectare	One veteran tree per hectare	No veteran trees present in woodland	2	2
Amount of deadwood	50% of all survey plots within the woodland parcel have standing deadwood, large dead branches/ stems and stumps	Between 25% and 50% of all survey plots within the woodland parcel have standing deadwood, large dead branches/ stems and stumps	Less than 25% of all survey plots within the woodland parcel have standing deadwood, large dead branches/ stems and stumps	2	2
Woodland disturbance	No nutrient enrichment or damaged ground evident	Less than 1 hectare in total of nutrient enrichment across woodland area and/or less than 20% of	More than 1 hectare of nutrient enrichment and/or more than 20% of	3	3

	Score			Habitat Parcel reference (see Figure A3.4)		
Indicator	Good (3 points)	Moderate (2 points)	Poor (1 point)	W1f7- 1	W1f7- 2	
		woodland area has damaged ground	woodland area has damaged ground			
TOTAL SCORE	-	-	-	35	34	

Habitat Quality	Assessment Criteria	Qualifying Habitat Parcels
Good	Scores > 32 (33 to 39)	W1f7-1 W1f7-2
Moderate	Scores 26 to 32	None
Poor	Scores < 26	None

3. Line of trees condition assessment

Habitat Description – Line of trees

A line of trees at least 20m in length with open habitat on each side.

	Habitat Parcel reference (see Figure A3.4)
Assessment Criteria	W1g6-2
More than 70% of trees are native species	✓
Tree canopy is predominantly continuous with gaps in canopy cover making up <10% of total area and no individual gap being >5m wide	\checkmark
Includes one or more mature or veteran tree	\checkmark
There is an undisturbed naturally vegetated strip of at least 6 m on both sides to protect the line of trees from farming and other anthropogenic operations	Х
At least 95% of the trees are in a healthy condition (excluding veteran features valuable for wildlife). There is little or no evidence of an adverse impact on tree health by damage from livestock or wild animals, pests or diseases, or human activity	✓

Habitat Quality	Assessment Criteria	Qualifying Habitat Parcels
Good	Passes 5 of 5 criteria	None
Moderate	Passes 3 or 4 of 5 criteria	W1g6-2
Poor	Passes 0, 1 or 2 of 5 criteria	None

4. Hedgerow condition

Attributes and functional groupings	Criteria	H2a-2	H2a-4	H2a-5	H2a-6	H2a-7	H2a-10	H2a-12*
A1. Height	>1.5 m average along length	✓	√	√	√	~	✓	-
A2. Width	>1.5 m average along length	✓	Х	\checkmark	\checkmark	\checkmark	\checkmark	-
B1. Gap – hedge base	Gap between ground and base of canopy <0.5m for >90% of length (unless 'line of trees')	✓	Х	✓	✓	Х	✓	-
B2. Gap - hedge canopy continuity	Gaps make up <10% of total length No canopy gaps >5 m	✓	√	\checkmark	✓	Х	√	-
C1. Undisturbed ground and perennial vegetation	 >1 m width of undisturbed ground with perennial herbaceous vegetation for >90% of length measured from outer edge of hedgerow, and is present on one side of the hedge (at least) 	✓	✓	✓	✓	•	✓	-
C2. Undesirable perennial vegetation	Plant species indicative of nutrient enrichment of soils dominate <20% cover of the area of undisturbed ground	✓	✓	~	~	✓	✓	-
D1. Invasive and neophyte species	>90% of the hedgerow and undisturbed ground is free of invasive non-native and neophyte species	✓	✓	✓	~	✓	✓	-
D2. Current damage	>90% of the hedgerow or undisturbed ground is free of damage caused by human activities	✓	✓	~	~	√	✓	-
	Additional group - applicable to	hedgero	ows with tro	ees only:				
E1. Tree age	At least one mature tree per 30m stretch of hedgerow. A mature tree is one that is	Х	N/A	✓	✓	Х	✓	-

Habitat Parcel reference (see Figure A3.4)

Habitat Parcel reference (see Figure A3.4)

Attributes and functional groupings	Criteria	H2a-2	H2a-4	H2a-5	H2a-6	H2a-7	H2a-10	H2a-12*
	at least 2/3 expected fully mature height for the species							
E2. Tree health	At least 95% of hedgerow trees are in a healthy condition (excluding veteran features valuable for wildlife). There is little or no evidence of an adverse impact on tree health by damage from livestock or wild animals, pests or diseases, or human activity	~	N/A	✓	✓	~	V	-

*As per limitation section – h2a-12 has not been conditioned assessed in the field and as a precautionary measure given a condition of good based of aerial mapping, and nearby hedgerows in the area.

Habitat Quality	Assessment Criteria	Qualifying Habitat Parcels
Condition c	ategories for hedgerows without trees:	
Good	No more than 2 failures in total; AND No more than 1 in any functional group	H2a-4
Moderate	No more than 4 failures in total; AND Does not fail both attributes in more than one functional group (e.g. fails attributes A1, A2, B1 & C2 = Moderate condition)	None
Poor	Fails a total of more than 4 attributes; OR Fails both attributes in more than one functional group (e.g. fails attributes A1, A2, B1 & B2 = Poor condition)	None
Condition c	ategories for hedgerows with trees:	
Good	No more than 2 failures in total; AND No more than 1 failure in any functional group	H2a-2 H2a-5 H2a-6 H2a-10 H2a-12
Moderate	No more than 5 failures in total; AND Does not fail both attributes in more than one functional group (e.g. fails attributes A1, A2, B1, C2 & E1 = Moderate condition)	H2a-7
Poor	Fails a total of more than 5 attributes; OR Fails both attributes in more than one functional group (e.g. fails attributes A1, A2, B1 & B2 = Poor condition)	None

Annex 4: Bat Survey Results

- Table 1 PGLRA and tree climbing bat survey results
- Table 2 Bat emergence/ re-entry meta data
- Table 3 Roost analysis

Tree ID	OS Grid Ref	Species & DBH (m)	Potential Roost Feature	Roost Potential suitability	Hibernation potential	Hibernation survey results	Active Season survey method	Roost
TC1	TL 84222 37186	Oak – 1.25	 PRF 1: Semi torn-out branch. Some cracks and rotting but nothing significant. Rotting bowl at base - low. PRF 2: Same as 1) – low. PRF 3: Trunk cavity at 6m, north-east facing. Entrance diameter is 8cm and feature extends diagonally upwards for 8cm. Feature also goes down 1cm. Internal diameter 10cm max., with dome-shaped apex. Woodlice inside with dirty, smooth and dry interior – moderate. PRF 4: Minor lifted bark – negligible. 	Moderate	No	N/A	Climbing inspection	No roost
TC2	TL 84267 37188	Ash – 0.75 (multi- stem)	 PRF 1: Knothole / callus roll located 8m high on main stem. Entrance 3cm leading inwards 10 cm then narrowing to cone. Internal diameter 3cm. Interior was rough, dry and fairly clean – low. PRF 2: Tear out feature - part of rotten heartwood, leaving small cavity. Feature leads 20cm inwards with the entrance measuring 2x3cm. Internal cavity was 3cm wide, dry, clean, smooth and sheltered. Good surrounding habitat – high. 	High	Yes	No change to PRFs from previous inspection, no bats recorded.	Climbing inspection	No roost
TC3	TL 84317 37195	Ash -1	 Ash -1 PRF 1: Entrance approx.7cm in diameter, extending 20cm inwards before tapering slightly. Rough and dirty inside with some decay forming a crevice that extends inwards by 5cm. Suitable for 1 or 2 bats - moderate. PRF 2: Both have entrances 7cm in diameter. Both lead inwards 20cm without tapering without leading up or down. Both have nesting material in and are dry, smooth, and clean. Both are exposed to predation - low. PRF 3: Entrance measures 7cm high and 3cm wide. Leads into gap behind rotting heartwood, leading 		Yes	Tree unsafe to climb – fungal brackets below feature	Climbing inspection	No roost

Table 1: Preliminary Ground Level Roost Assessment and tree climbing bat survey results

Tree ID	OS Grid Ref	Species & DBH (m)	Potential Roost Feature	Roost Potential suitability	Hibernation potential	Hibernation survey results	Active Season survey method	Roost
			10cm up and 7cm inwards. Damp and rough inside with no interspecies competition – moderate.					
TC4	TL 84344 37166	Oak -0.9	PRF 1: Small cavity in wound growth around limb tear out. Facing south-west, 10m high.PRF 2: Two cavities on underside of northern aspect of tree Approx.16-18m high.	Moderate	No	N/A	Climbing inspection	No roost
TC5	TL 84333 37166	Oak – 0.75	No suitable space/ insulation for bats following aerial inspection – negligible.	Negligible	No	N/A	Climbing inspection	No roost
TC6	TL 84357 37145	Oak – 0.9	PRF 1: Wound growth around limb tear out potentially leading to cavity behind. Feature at 7m facing west.	Moderate	No	N/A	Climbing inspection	No roost
TC7	TL 84359 37143	Oak -1.1	PRF 1: Crack on underside of torn branch on northern side of tree. South-west facing, approx.10m high.	Moderate	No	N/A	Climbing inspection	No roost
TC8	TL 84367 37160	Oak -0.8	PRF 1: Wound growth around limb tear out potentially leading to cavity. Feature at 10m facing north-west.	Moderate	No	N/A	Climbing inspection	No roost
TC9	TL 84398 37168	Oak -0.9	PRF 1: Rot hole at 7m facing north-east with dead, cracked branch 1m above.	Moderate	No	N/A	Climbing inspection	No roost
TC10	TL 84414 37148	Oak -0.8	PRF 1: Branch cavity on fork, approx. 4m high facing south-east.	Moderate	Yes	No change to PRFs from previous inspection, no bats recorded.	Climbing inspection	No roost
TC11	TL 84486 37164	Oak -0.9	PRF 1: Rot hole approx. 9m high facing south. Observed from south field.	Moderate	No	N/A	Climbing inspection	No roost

Tree ID	OS Grid Ref	Species & DBH (m)	Potential Roost Feature	Roost Potential suitability	Hibernation potential	Hibernation survey results	Active Season survey method	Roost
TC12	TL 84512 37214	Oak -1	 PRF 1: Tear out and cracks at base of dead branch at approx. 6m, facing south. Wounds are very open with no crevices. Damp and dirty inside with small amount of lifted bark – low. PRF 2: Lifted bark open and exposed. Gap between bark and branch is 4cm. Extends 60cm along branch. Damp and dirty inside – low. 	Low	No	N/A	Climbing inspection	No roost
TC13	TL 84557 37199	Oak -1.25	 PRF 1: Feature measures15x3cm, heartwood is desiccated with cavity inside leading in 10cm. Cavity is quite sheltered, dry and suitable for a single bat – moderate. PRF 2: Tightly sealed, small cavity measuring 1x2cm. Full of woodlice and facing south – negative. PRF 3: Branch cavity at approx. 8m facing southwest. Entrance measures 12x4cm with internal 2cm cavity. Interior is dirty and full of debris. Leads horizontally towards stem for 10cm – low. 	Moderate	No	N/A	Climbing inspection	No roost
TC14	TL 84581 37197	Oak -1	 PRF 1: Tear out is approx.1m high and 40cm wide. Cavity at the top goes in 10cm with entrance 3x7cm. Internal is similar dimensions. Sheltered, dry, dirty – moderate. PRF 2: Knot hole with peg on branch at approx. 6m extending over field to the south. Opens into crack running along top of branch. Crack is 3cm wide and 3cm deep. Open and exposed all the way along – low. 	Moderate	Yes	No change to PRFs from previous inspection, no bats recorded.	Climbing inspection	No roost
TC15	TL 84599 37211	Ash -1.1	 PRF 1: Entrance faces downward and is 6cm in diameter. Internal diameter is 6cm and leads towards stem. Clean and smooth interior leading inwards 15cm No competitors – high. PRF 2: Entrance measures 3x3cm in diameter leading into longer cavity. Extends 20cm into trunk. Fairly damp and dirty interior but suitable space - moderate. 	High	Yes	No change to PRFs from previous inspection, no bats recorded.	Climbing inspection	No roost

Tree ID	OS Grid Ref	Species & DBH (m)	Potential Roost Feature	Roost Potential suitability	Hibernation potential	Hibernation survey results	Active Season survey method	Roost
			PRF 3: Open and exposed with some crevices but no suitable space or insulation – low.					
TC16	TL 84630 37199	Oak -1.2	 PRF 1: Entrance 5cm diameter leading back 20cm. Internal diameter 5cm. Interior is dry with low levels of dirt, cobwebs and woodlice. Rough and bumpy interior - moderate. PRF 2: Entrance measures 10x4c with internal diameter of 5cm. Leads back 15cm and is dry, dirty, full of debris and cobwebs – moderate. 	Moderate	Yes	No change to PRFs from previous inspection, no bats recorded.	Climbing inspection	No roost
TC17	TL 84652 37164	Oak -1	PRF 1: Open and exposed. Entrance 12x12cm leading back 25cm – low.	Low	No	N/A	N/A	No roost
TC19	TL 84457 37025	Oak -0.75	PRF 1: Tear out with heartwood in wound. Cavity in heartwood with top entrance measuring 5cm in diameter. Goes diagonally upwards and interior is clean, dry, smooth with a few woodlice. Feature narrows and goes up 40cm– high. PRF 2: Second entrance 50cm lower facing north. Entrance measures 5x3cm and narrows to dome apex 20cm up. Internal diameter measures 6cm, is sheltered and has few cobwebs – high.	High	Yes	No change to PRFs from previous inspection, no bats recorded.	Climbing inspection	No roost
TC20	TL 84405 37036	Oak -0.7	PRF 1: Superficial only – does not lead anywhere – negligible PRF 2: Superficial only – does not lead anywhere – negligible	Negligible	N/A	N/A	Climbing inspection	No roost
TC21	TL 84371 37026	Oak -0.8	PRF 1: Tear out is 2.5m long and 30cm wide. Calluses around edges do not have roosting potential. Small vertical cracks in exposed heartwood are superficial. Top of feature is open and exposed. Rough, clean and dry interior – low.	Low	No	N/A	N/A	No roost

Tree ID	OS Grid Ref	Species & DBH (m)	Potential Roost Feature	Roost Potential suitability	Hibernation potential	Hibernation survey results	Active Season survey method	Roost
TC22	TL 84330 37042	Oak -0.9	PRF 1: Knot hole10x4cm is completely open. Goes inwards by 5cm and is exposed – low. PRF 2: More of a bird nesting feature - no animals present. Entrance 15x10cm, secondary entrance 5cm diameter. Bowl inside - too open for bats. Goes in 30cm. straw at base with few cobwebs. Cavity goes up 10cm and is suitable for a small number of bats – moderate.	Moderate	No	N/A	Climbing inspection	No roost
TC23	TL 84310 37044	Oak -0.5	 PRF 1: Entrance measures 4x10cm at bottom of feature. Goes up 35cm inside. Very dirty and rough inside. Internal width 12cm and depth 3cm – moderate. PRF 2: Second cavity at top of heartwood panel. Entrance measures 10x4cm. Goes up into main trunk of tree by 60cm. Internal diameter measures 6cm all the way up. Dry, fairly clean with no competitors – high. 	High	Yes	No change to PRFs from previous inspection, no bats recorded.	Climbing inspection	No roost
TC24	TL 84289 37036	Oak -1.1	PRF 1: Wound is 60x15cm. No crevices behind exposed heartwood. Cavity at top left of feature which has a 3cm diameter entrance and goes in 6cm. Sheltered, dry and clean interior – moderate.	Moderate	No	N/A	Climbing inspection	No roost
TC25	TL 84259 37032	Ash -0.6	 PRF 1: Entrance 40cm high, 5cm wide. Cavity on southern side extends upwards into southern stem. Internal diameter 4cm goes up 30cm. Damp inside, secure and sheltered, smooth and mostly clean - high. PRF 2: Knot hole facing south. Entrance 6cm in diameter extending 25cm inside trunk. Interior damp with some sludge but no competitors – moderate. 	High	Yes	No change to PRFs from previous inspection, no bats recorded.	Climbing inspection	No roost
TC26	TL 84259 37028	Oak -0.5	PRF 1: Entrance measures 6cm in diameter and leads inwards 10cm before tapering slightly. Doesn't go up or down with interior rough and dry – low.	Low	No	N/A	N/A	No roost

Tree ID	OS Grid Ref	Species & DBH (m)	Potential Roost Feature	Roost Potential suitability	Hibernation potential	Hibernation survey results	Active Season survey method	Roost
TC27	TL 84257 36963	Oak -0.8	 PRF 1: Rim of entrance smooth possibly from mammal use. Bowl shaped cavity full of sludge and slugs. Damp, wet and dirty interior. Entrance measures 7cm in diameter extending back 20cm. Internal diameter measures 10cm – low. PRF 2: Ants nest inside. Crevice at back of wound is 7x2cm leading inwards 7cm. Dry and sheltered with lots of debris and dirt – moderate. PRF 3: Entrance measures 15x12cm leading inwards 20cm and diagonally downwards. Rough and dirty interior with peg of old branch sticking out – low. 	Confirmed – 2-3 myotis found with endo	Yes	No change to PRFs from previous inspection, no bats recorded.	Climbing inspection	Roost identified – 2-3 Natterer's bat (Myotis nattereri) confirmed by DNA of droppings – day roost
TC28	TL 84274 36944	Oak -0.9	 PRF 1: No suitable space or insulation – negative. PRF 2: Crevices and deadwood in limb with loose bark at same height as callous roll. Main part is hollowed out. Cavity entrance is 5x4cm leading inwards 10cm into a wedge leading upwards by 10cm. Clean, sheltered and dry interior with some debris - moderate. PRF 3: Branch facing east with cavity on top extending along branch 5m. Entrance not visible, but measures 20x10cm. Internal diameter measures 9cm. Interior is clean and smooth with debris at entrance before tapering tapers to end (approx. 70 cm) – high. 	High	Yes	No change to PRFs from previous inspection, no bats recorded.	Climbing inspection	No roost
TC29	TL 84318 36817	Ash -0.6	PRF 1: Woodpecker hole at top of wound feature at 9m, on eastern aspect of main stem. Entrance measures 7cm and opens out into cavern in centre of tree. Internal diameter widens to 10cm. It goes back 20cm. Bird nesting material in base. Rough, dry, clean with woodlice. Two crevices at back on top could be suitable for a bat – moderate. PRF 2: Callus roll on main stem, no potential – negative.	Moderate	Yes	Squirrels present. No change to PRFs from previous inspection, no bats recorded.	Climbing inspection	No roost

Tree ID	OS Grid Ref	Species & DBH (m)	Potential Roost Feature	Roost Potential suitability	Hibernation potential	Hibernation survey results	Active Season survey method	Roost
TC32	TL 84260 36974	Oak -0.8	 PRF 1: Woodpecker hole at 4m, facing north on main stem. Entrance measures 8cm in diameter. Opens out into large cavity. Bird nesting material in base (old). Goes back 30cm. Internal diameter 15cm. Rough, clean, dry. Goes up 20cm, tapering to spire. Dry, sheltered – high. PRF 2: Woodpecker hole at 6m, facing north-west on main stem. Entrance measures 5cm in diameter and goes in 15cm. Internal diameter measures 6cm. Dry, dusty, cobwebs. Goes up 2cm – moderate. 	High	Yes	No change to PRFs from previous inspection, no bats recorded.		No roost
TC33	TL 84259 37034	Oak -0.5	PRF 1: Knot hole at 4m on main stem facing north west Entrance measures 4cm wide and goes in 8cm. Feature doesn't extend down. Crevices at top and right that go up 5cm. Clean, dry, sheltered. Room for 2 or 3 bats. Uncluttered drop zone – moderate.	Moderate	No	N/A	Climbing inspection	No roost
T1	TL 84267 37214	Ash - 1.2	PRF 1: Cavity tapers to spire at apex. Internal diameter 25cm. Damp inside with debris, fresh cobwebs and bobbly sides. Reasonably clean with some debris.	High	Yes	No change to PRFs from previous inspection, no bats recorded.	Emergence/re- entry	No roost
T2	TL 84322 37179	Ash -0.7	PRF 1: Open trunk cavity (frost crack) running from ground level up to 6m where it seems to extend into sheltered internal cavity for at least 20cm.	N/A	Yes	No change to PRFs from previous inspection, no bats recorded.	Emergence/re- entry	No roost
Т3	TL 84566 37200	Oak (dead tree) -0.8	PRF 1: Multiple woodpecker holes facing south-west at 5m.	N/A	No	N/A	Emergence/re- entry	No roost
Τ4	TL 84555 37029	Ash -1.5 (multi- stem)	PRF1: 4 stems, eastern stem goes up extensively and so does the northern stem. Eastern and northern stem goes up at least 1 m, some debris and dirty and damp, narrows as it goes up. Internal dimensions 20 cm at base.	High	Yes	No change to PRFs from previous inspection, no bats recorded.	Emergence/re- entry	No roost

Tree ID	OS Grid Ref	Species & DBH (m)	Potential Roost Feature	Roost Potential suitability	Hibernation potential	Hibernation survey results	Active Season survey method	Roost
Τ5	TL 84498 37026	Ash -0.8	PRF 1: Splits and raised bark on all aspects starting from 1m. Diseased tree.	N/A	No	N/A	Emergence/re- entry	No roost
Τ6	TL 84448 37017	Likely oak (dead tree) -0.8	PRF 1: Loose bark on all aspects. PRF 2: Large opening on tree facing east.	N/A	Yes	Unsafe to climb – bark torched from ground level. No change to PRFs from previous inspection, no bats recorded.	Emergence/re- entry	No roost
Τ7	TL 84370 37017	 TL Likely oak PRF 1: Entrance measures 30x10cm. Open cavity 84370 (dead with some shelter behind callous rolls - low. 37017 tree) - PRF 2: Cavity on west facing limb at 9m. Knot hole leads to crevice at back. Entrance measures 10x7cm and extends back 6cm and down by 2cm. Interior is rough, dry, open and with debris – low. PRF 3: Central dead stem at 8m with multiple entrances on all aspects. Hollow inside but open and exposed at top - low. PRF 4: Knot hole at 7.5m facing south-east. Entrance measures 8x6cm and has a chimney that extends up by10cm. Interior is sheltered and dry – moderate. PRF 5: Wound in upright limb at 7m facing south-east. Feature extends 10x8cm and goes up by 40 cm. Interior is rough, clean, dry and with cobwebs. Extends down by 2cm, with some decay at base - high. PRF 6: Knothole entrance measuring 9x9cm, approx. 5m high and facing south-east. Internal diameter measures 15cm and goes diagonally upwards by 80cm. Interior is smooth, clean and dry – high. 		High	Yes	Unsafe to climb – bark torched from ground level. No change to PRFs from previous inspection, no bats recorded.	Emergence/re- entry	No roost

Tree ID	OS Grid Ref	Species & DBH (m)	Potential Roost Feature	Roost Potential suitability	Hibernation potential	Hibernation survey results	Active Season survey method	Roost
			diameter measures 15cm before narrowing. Multiple chambers at top, clean and dry – high.					
Т8	TL 84309 37039	Oak -0.6	PRF1: Disease in trunk leading to numerous large cavities from 2m up to 7m facing west. PRF2: Rot hole facing north-west at 7m.	N/A	Yes	Unsafe to climb	Emergence/re- entry	No roost
Τ9	TL 84273 37002	Oak -0.6	PRF1: Large hollow base, dry inside, smooth, goes upwards 2 m. The top 50 cm is dirty and damp. The lower part is dry. Entrance and internal - 20 cm diameter.	High	Yes	Unsafe to climb –inspection from ground level. No change to PRFs from previous inspection, no bats recorded.	Emergence/re- entry	No roost
T10	TL 84280 36910	Likely oak (dead tree) -1	PRF1: Dead tree with loose bark on trunk and branches - starting from 2m facing west.	N/A	Yes	Unsafe to climb –inspected from ground level. No change to PRFs from previous inspection, no bats recorded.	Emergence/re- entry	No roost
T11	TL 84260 36909	Ash -0.5	PRF1: Woodpecker hole at 12m on eastern side. PRF2: Cavity in branch on south-western side at 12m.	N/A	No	N/A	Emergence/re- entry	No roost
T12	TL 84287 36835	Oak -1.1	PRF1: Callus roll in dead branch facing west.	N/A	No	N/A	Emergence/re- entry	No roost
T13	TL 84296 36824	Oak -1	PRF1: Dead, rotten tree with numerous woodpecker and rot holes facing north at 5m. Loose bark on all aspects.	N/A	Yes	Unsafe to climb –inspected from ground level. No change to PRFs from previous	Emergence/re- entry	No roost

Tree ID	OS Grid Ref	Species & DBH (m)	Potential Roost Feature	Roost Potential suitability	Hibernation Hibernation potential survey resu		Active Season survey method	Roost
						inspection, no bats recorded.		
T15	TL 84315 36792	Aspen - 0.4	PRF1: Three woodpecker holes on western side of tree and approx. 10-12m.	N/A	No	N/A	Emergence/re- entry	No roost

Date	Trees Surveyed	Visit Number	Survey type	Sunset/sunrise time	Survey start time	Survey end time	Temperature (°C)	Rain (1-4)	Wind (Beaufort)	Cloud cover (0-8)
04-08-21	T4, T7-9	1/3	Dawn	05:23	03:23	05:38	9-11	1	0	0
05-08-21	T1-2	T1, T2 - 1/3	Dawn	05:24	03:23	05:39	11-12	1	0	5
24-08-21	T6-9	T6 – 1/2 T7 – 2/3 T8, T9 – 2/3	Dawn	05:55	03:55	06:10	12	1	1	6
25-08-21	T1-5	T1, T2, T4 – 2/3 T3, T5 – 1/2	Dusk	19:59	19:20	19:44	16	1	2	7
26-08-21	T10-11	1/2	Dusk	19:57	19:42	21:57	14	1	3	8
21-09-21	T1-3	T1, T2 – 3/3 T3 – 2/2	Dawn	06:41	04:41	06:56	13-14	1	1	4
22-09-21	T10-11	2/2	Dawn	06:42	04:42	06:57	11-14	1	1	0
28-09-21	T4-5	T4 – 3/3 T5 – 2/2	Dawn	06:53	04:53	07:08	10	1	1	2

Table 2: Emergence and re-entry survey metadata



Table 3: Roost analysis of bat recorded during dusk survey on T16 – sound call

Annex 5 Photos of Habitats and Ecological Features

Woodland Habitat – Ancient and Broadleaved



Photograph 1: Butler's Wood ancient woodland adjacent to A131 (Parcel no. w1f7-1).



Photograph 2: Butler's Wood ancient woodland adjacent to A131 (Parcel no. w1f7-1).



Photograph 3: Butler's Wood ancient woodland adjacent to A131 (Parcel no. w1f7-1).



Photograph 5: Western edge of Waldegrave Wood ancient woodland (Parcel no. w1f7-2).



Photograph 4: Northern edge of Waldegrave Wood ancient woodland (Parcel no. w1f7-2).



Photograph 6: Waldegrave Wood ancient woodland understory composition (Parcel no. w1f7-2).



Photograph 7: Waldegrave Wood ancient woodland understory composition (Parcel no. w1f7-2).



Photograph 8: Broadleaved woodland around Pond 7, east of A131 (Parcel no. w1g7-1).



Photograph 9: Mature multi-stem willow tree within broadleaved woodland surrounding Pond 7, east of A131 (Parcel no. w1g7-1).



Photograph 10: Broadleaved woodland adjacent to Pond 18 located south of the scheme (Parcel no. w1g7-2).

Grassland Habitat – Amenity, Neutral and Grazed



Photograph 11: Grazed grassland (Parcel no. g3c-1).



Photograph 12: Grass Margin adjacent to arable field (Parcel no. g3c-3).



Photograph 13: Grass Margin running parallel to line of trees (Parcel no. g3c-8).



Photograph 14: Grassland road verge (Parcel no. g3c-6).



Photograph 15: Neutral Grassland (Parcel no. g3c-4).



Photograph 16: Amenity Grassland (Parcel no. g4-2).

Field Boundaries - Hedgerows and Lines of Trees



Photograph 17: Line of planted poplar trees south of Old Road and adjacent to intensive orchards (Parcel no: w1g6-6)



Photograph 18: Hedgerow with trees north of Old Road (Parcel no: h2a-1)





Photograph 19: Line of trees adjacent to track north of Lodge Farm. (Parcel no: w1g6-11)

Photograph 20: Line of trees/mature hedgerow running along the edge of the Scout Campgrounds, east of A131. (Parcel no: w1g6-11)



Photograph 21: Hedgerow along Old Road just east of A131 and south of Waldegrave Wood. (Parcel no: h2a-5 and H2)



Photograph 22: Line of Lime Trees in the southwestern corner of the scheme leading to Stone's Farm. (Parcel no: w1g6-4 & w1g6-5)



Ponds – Positive GCN Presence



Photograph 24: P3 situated in the north-west corner of Butler's Wood.



Photograph 26: P5 situated on the southern side of Butler's Wood.



Photograph 25: P4 situated in the north-west corner of Butler's Wood.



Photograph 27: P6 situated on the southern side of Butler's Wood.



Photograph 28: P14 situated on Old Road just north of Nether House Farm.



Photograph 29: P20 situated in the southwestern corner of the scheme.

Ecological Constraints – Bats, Badgers and Invasive Species



Photograph 30: T16 Confirmed Bat Roost located in the south-western corner of Waldegrave Wood.



Photograph 32: Tree C27 Confirmed Bat Roost located in Waldegrave Wood.



Photograph 34: Outlier badger sett entrance located along western edge of Butler's Wood.



Photograph 31: Infra-Red Camera Footage of Soprano Pipistrelle perching outside bat roost feature on T16.



Photograph 33: Active badger sett entrance located along southern edge of Butler's Wood.



Photograph 35: Mammal path leading to outlier badger sett located along western edge of Butler's Wood.



Photograph 36: Disused outlier badger sett. Two entrances located within small broadleaved woodland surrounding Pond 7.



Photograph 38: Annex/Subsidiary badger sett entrance facing SE within pile of debris located on Scout Campgrounds.



Photograph 40: Main badger sett with 18 active entrances located along a hedgerow north-east of Lodge Farm.



Photograph 37: Two badger latrines located within Scout Campgrounds east of A131.



Photograph 39: One of three entrances to outlier badger sett located along arable field margin north-east of Scout campgrounds.



Photograph 41: Outlier badger sett entrance at base of a mature oak tree located along a hedgerow north-east of Lodge Farm.


Photograph 42: Main badger sett with 10 entrances located in line of trees on bank situated east of Nether House Farm.



Photograph 43: Main badger sett with 10 entrances located in line of trees on bank situated east of Nether House Farm.



Photo 44: Variegated Yellow Archangel located within Waldegrave Wood



Photo 45: Variegated Yellow Archangel located within grassland surrounding Scout Campgrounds

National Grid plc National Grid House Warwick Technology Park Gallows Hill, Warwick CV34 6DA United Kingdom

Registered in England and Wales No. 4031152

nationalgrid.com